Universität Stuttgart

Fachbereich Mathematik

Iterated fibre sums of algebraic Lefschetz fibrations M. J. D. Hamilton

Preprint 2012/009

Universität Stuttgart

Fachbereich Mathematik

Iterated fibre sums of algebraic Lefschetz fibrations M. J. D. Hamilton

Preprint 2012/009

Fachbereich Mathematik Fakultät Mathematik und Physik Universität Stuttgart Pfaffenwaldring 57 D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de
WWW: http://www.mathematik.uni-stuttgart.de/preprints

ISSN 1613-8309

@ Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors. $\mbox{\it Le}T_{E}X\mbox{-}Style:$ Winfried Geis, Thomas Merkle

ITERATED FIBRE SUMS OF ALGEBRAIC LEFSCHETZ FIBRATIONS

ABSTRACT. Let M denote the total space of a Lefschetz fibration, obtained by blowing up a Lefschetz pencil on an algebraic surface. We consider the *n*-fold fibre sum M(n), generalizing the construction of the elliptic surfaces E(n). For a Lefschetz pencil on a simply-connected minimal surface of general type we partially calculate the Seiberg-Witten invariants of the fibre sum M(n) using a formula of Morgan-Szabó-Taubes. As an application we derive an obstruction for self-diffeomorphisms of the boundary of the tubular neighbourhood of a general fibre in M(n) to extend over the complement of the neighbourhood. Similar obstructions are known in the case of elliptic surfaces.

1. INTRODUCTION

It is well-known that the sequence of simply-connected elliptic surfaces E(n) without multiple fibres can be constructed from the elliptic surface E(1), diffeomorphic to $\mathbb{CP}^2 \# 9\overline{\mathbb{CP}}^2$, by fibre summing along general fibres [6]. The elliptic fibration on E(1) can be obtained by considering a certain Veronese embedding of \mathbb{CP}^2 into a complex projective space and then taking the blow-up of a Lefschetz pencil. We consider the following generalization: Let M' denote an arbitrary smooth algebraic surface. It admits a Lefschetz pencil which extends to a Lefschetz fibration on some blow-up M. We can consider the iterated fibre sums of these fibrations, yielding a sequence M(n) of symplectic manifolds with an induced Lefschetz fibration over \mathbb{CP}^1 .

We describe the basic topology of these manifolds in Section 4. For example, if M is simplyconnected, then all M(n) are simply-connected and there is a description of the intersection form and of the canonical class. Considering the Seiberg-Witten invariants of M(n) we show that if M' is a minimal surface of general type, then the only basic class up to sign of M(n) for all $n \ge 2$ which has non-zero intersection with the fibre is the canonical class. To determine the Seiberg-Witten basic classes of the fibre sum M(n) we use a formula of Morgan, Szabó and Taubes. In general this formula involves a sum over several characteristic classes. However, in the case of the manifolds M(n) there is only one summand and the formula completely determines the basic classes under the mentioned constraint.

We then give an application of these calculations to the question which orientation preserving self-diffeomorphisms of the boundary of the tubular neighbourhood $\nu\Sigma$ of a general fibre in M(n) extend over the complement $M(n) \setminus \operatorname{int} \nu\Sigma$. In the case of elliptic surfaces E(n) a complete answer to the corresponding question is known [6]: For E(1) every diffeomorphism on $\partial\nu\Sigma$ extends over $E(1) \setminus \operatorname{int} \nu\Sigma$ and for E(n) with $n \ge 2$ only if it preserves the torus fibration on the boundary. For the manifolds M(n) we derive a criterion that can be used to show that certain diffeomorphisms do not extend over the complement.

2. LEFSCHETZ FIBRATIONS

Let M' be a smooth algebraic surface in some \mathbb{CP}^N of degree r so that

$$[M'] = r[\mathbb{CP}^2] \in H_4(\mathbb{CP}^N; \mathbb{Z})$$

Date: November 6, 2012.

²⁰¹⁰ Mathematics Subject Classification. Primary 57R57, 57R17; Secondary 14J29. *Key words and phrases.* 4-manifold, symplectic, fibre sum, Lefschetz fibration.

Choose a hyperplane $H \cong \mathbb{CP}^{N-1}$ in \mathbb{CP}^N intersecting M' transversely. According to Section 1.5 and 1.6 in [10] there exists a Lefschetz pencil containing H as one of its hyperplanes. A pencil is the set of hyperplanes which contain a given linear subspace $A \cong \mathbb{CP}^{N-2}$, called the axis of the pencil. The axis A intersects M' transversely in r points forming the base locus B. Blowing up these points results in a fibration

$$M = M' \# r \overline{\mathbb{CP}^2} \longrightarrow \mathbb{CP}^1.$$

This is what we call an *algebraic Lefschetz fibration*. A Lefschetz fibration has finitely many singular fibres where the singularities have a certain normal form. We can assume that each singular fibre has precisely one singularity. The generic fibre Σ_M is given by the proper transform of a non-singular hyperplane section $\Sigma_{M'}$ in M' determined by a generic hyperplane in the pencil. We denote the genus of Σ_M by g.

By the first Lefschetz Hyperplane theorem, the homomorphism

$$i_M \colon H_1(\Sigma_M; \mathbb{Z}) \to H_1(M; \mathbb{Z}),$$

induced by inclusion for a smooth fibre Σ_M is a surjection. The so-called second Lefschetz Hyperplane theorem [1] shows that the kernel of this map is generated by the set of *vanishing cycles*. The vanishing cycles bound embedded disks in M, called *Lefschetz thimbles* or *vanishing disks*, which intersect Σ_M only in the vanishing cycle and contain precisely one critical point of the fibration. The vanishing disks are formed above certain arcs in \mathbb{CP}^1 . For each critical point there is a corresponding vanishing cycle and a vanishing disk. If we frame a vanishing disk D on its boundary by the direction normal to the cycle inside the fibre, then D has self-intersection equal to -1.

The fibration defines a natural framing for the tubular neighbourhood of a general fibre Σ_M in M. Consider a diffeomorphism between the fibres in two copies of M that identifies the vanishing cycles. Lift the diffeomorphism in the standard way to an orientation reversing diffeomorphism of the boundary of the tubular neighbourhoods using the framing determined by the fibration. If we form the generalized fibre sum, we get a closed symplectic 4-manifold

$$M(2) = M \#_{\Sigma_M = \Sigma_M} M.$$

We can iterate the construction to get symplectic 4-manifolds M(n), where

$$M(n) = M \#_{\Sigma_M = \Sigma_M} M \#_{\Sigma_M = \Sigma_M} \dots \#_{\Sigma_M = \Sigma_M} M.$$

By our choice of gluing the Lefschetz fibration on M extends to a symplectic Lefschetz fibration on M(n).

3. FIBRE SUMS AND THE MORGAN-SZABÓ-TAUBES FORMULA

We recall some results about generalized fibre sums [7, 8]. Let M and N denote closed oriented 4-manifolds with closed oriented embedded surfaces Σ_M and Σ_N of genus g and self-intersection zero. We choose embeddings

$$i_M: \Sigma \to M$$
$$i_N: \Sigma \to N$$

that realize the surfaces as images of a closed oriented surface Σ . We fix framings $\Sigma \times D^2$ of the closed tubular neighbourhoods $\nu \Sigma_M$ and $\nu \Sigma_N$ and denote the manifolds minus the interior of the neighbourhoods by M_0 and N_0 . We want to glue M_0 and N_0 together using an orientation reversing diffeomorphism $\phi : \partial M_0 \to \partial N_0$ that preserves the circle fibration and covers the diffeomorphism

 $i_N \circ i_M^{-1}$ between the surfaces. In the chosen framings any such diffeomorphism is isotopic to a diffeomorphism of the form

$$\phi: \Sigma \times S^1 \longrightarrow \Sigma \times S^1,$$
$$(x, \alpha) \mapsto (x, C(x) \cdot \overline{\alpha})$$

where the bar denotes complex conjugation and $C : \Sigma \to S^1$ is a smooth map. The map ϕ depends up to isotopy only on the cohomology class $C^* d\alpha \in H^1(\Sigma; \mathbb{Z})$ that we also denote by C. The result of the generalized fibre sum is

$$X = M \#_{\Sigma_M = \Sigma_N} N = M_0 \cup_{\phi} N_0$$

and in general depends on the choice of the cohomology class C.

It is well-known that the Euler characteristic and the signature of X are given by

$$e(X) = e(M) + e(N) + 4g - \sigma(X) = \sigma(M) + \sigma(N).$$

4

Suppose that Σ_M and Σ_N represent indivisible homology classes. Then the first homology of X is given by the cokernel of the homomorphism

$$i_M \oplus i_N : H_1(\Sigma; \mathbb{Z}) \to H_1(M; \mathbb{Z}) \oplus H_1(N; \mathbb{Z})$$

induced by the embeddings.

We want to describe the second homology of X. Assume that M, N and X have torsion free homology and Σ_M and Σ_N represent indivisible classes. The framings determine push-offs of Σ_M and Σ_N into the boundaries ∂M_0 and ∂N_0 . Under inclusion in X we get surfaces Σ_X and Σ'_X . There also exist surfaces B_M and B_N in M and N which intersect Σ_M and Σ_N in a single positive transverse point. Since the gluing preserves the meridians $\{*\} \times S^1$ these surfaces sew together in X to a surface B_X . We define P(M) to be the orthogonal complement of the span of Σ_M and B_M in $H_2(M;\mathbb{Z})$, and similarly for N. A curve on Σ_M times the meridian σ_M defines a torus on ∂M_0 . Under inclusion in X this is a torus of self-intersection zero, called a *rim torus*. It is null-homologous in M but not necessarily in X. There are also so called *vanishing surfaces* in X (see [4]), sewed together along curves on the push-off of Σ_M in ∂M_0 and a curve on ∂N_0 which get identified under gluing and bound in M_0 and N_0 . With these preparations we can describe the second homology of X and the intersection form. Let c denote the rank of the kernel of the homomorphism $i_M \oplus i_N$ above.

Theorem 1. There exists a basis S_1, \ldots, S_c of the vanishing classes S'(X) and a basis R_1, \ldots, R_c of the rim tori R(X) such that there exists a splitting

$$H_2(X;\mathbb{Z}) = P(M) \oplus P(N) \oplus (S'(X) \oplus R(X)) \oplus (\mathbb{Z}B_X \oplus \mathbb{Z}\Sigma_X).$$

where

$$(S'(X) \oplus R(X)) = (\mathbb{Z}S_1 \oplus \mathbb{Z}R_1) \oplus \ldots \oplus (\mathbb{Z}S_c \oplus \mathbb{Z}R_c).$$

The direct sums are all orthogonal, except the direct sums inside the brackets. In this decomposition of $H_2(X;\mathbb{Z})$, the restriction of the intersection form Q_X to P(M) and P(N) is equal to the intersection form induced from M and N and has the structure

$$\left(\begin{array}{cc} B_M^2 + B_N^2 & 1\\ 1 & 0 \end{array}\right)$$

on $\mathbb{Z}B_X \oplus \mathbb{Z}\Sigma_X$ and the structure

3

 $\left(\begin{array}{cc}S_i^2 & 1\\ 1 & 0\end{array}\right)$

on each summand $\mathbb{Z}S_i \oplus \mathbb{Z}R_i$. We call such a basis for $H_2(X;\mathbb{Z})$ a normal form basis.

The vanishing surfaces S_i are obtained from a basis $\alpha_1, \ldots, \alpha_c$ for the kernel of $i_M \oplus i_N$. Suppose we only have a generating set for this subgroup. Expressing each basis element α_i in terms of the generating set we have:

Proposition 2. Let $\tilde{S}_1, \ldots, \tilde{S}_e$ denote vanishing surfaces obtained from a generating set for the kernel of $i_M \oplus i_N$. Then we can write each vanishing surface S_i in Theorem 1 as a linear combination of the \tilde{S}_i and certain rim tori.

The rim tori are needed in this proposition to separate the different vanishing surfaces that we get. Generally speaking, we fix a basis of rim tori and then determine a dual basis of vanishing surfaces. A basis of rim tori in one generalized fibre sum determines such a basis in all of them.

There is a formula for the Seiberg-Witten invariants of X due to Morgan, Szabó and Taubes [13] that we want to describe in our notation. The formula works only if $g \ge 2$. Recall that the Seiberg-Witten invariant of a closed, oriented 4-manifold with $b_2^+ > 1$ is a map

$$SW_X: Spin^c(X) \to \mathbb{Z}.$$

There is a related invariant

$$SW_X: \mathcal{C}(X) \to \mathbb{Z}$$

defined on the set C(X) of characteristic elements in $H^2(X; \mathbb{Z})$ by summing over all $Spin^c$ -structures with the same first Chern class. Under our assumptions a $Spin^c$ -structure is determined by its Chern class. A characteristic class $k \in C(X)$ is called a Seiberg-Witten basic class if $SW_X(k) \neq 0$. Let k' be a Seiberg-Witten basic class of X. The adjunction inequality shows that k' has zero intersection with each rim torus, hence the Poincaré dual of k' has no vanishing surface component. The adjunction inequality also shows that $|k' \cdot \Sigma_X| \leq 2g - 2$. The formula only makes a statement about the case that $k' \cdot \Sigma_X = \pm (2g - 2)$. We can restrict to the positive case: Let k denote a characteristic class on Xsuch that the Poincaré dual PD(k) is of the form

$$PD(k) = p_M + p_N + \sum_{i=1}^{c} \epsilon_i R_i + (2g - 2)B_X + \beta_X \Sigma_X,$$

with $p_M \in P(M)$ and $p_N \in P(N)$. One can show that

$$H^2(M_0;\mathbb{Z}) \cong H_2(M_0,\partial M_0;\mathbb{Z}) = P(M) \oplus \mathbb{Z}B'_M \oplus \ker i_M$$

where B'_M is the surface B_M with a disk deleted and i_M denotes the map on $H_1(\Sigma; \mathbb{Z})$ induced by inclusion. There are obvious restriction maps of $H^2(X; \mathbb{Z})$ and $H^2(M; \mathbb{Z})$ to $H^2(M_0; \mathbb{Z})$, and similarly for N. It follows that the set of characteristic classes on X which have the same square and the same restriction to M_0 and N_0 as k is given by

$$\mathcal{K}(k) = \{ l \in \mathcal{C}(X) \mid PD(l) = PD(k) + R \text{ with } R \in R(X) \}.$$

We also set

$$\mathcal{K}_M(k) = \{ l \in \mathcal{C}(M) \mid PD(l) = p_M + (2g - 2)B_M + \beta_M \Sigma_M \text{ with } \beta_M \in \mathbb{Z} \}$$

$$\mathcal{K}_N(k) = \{ l \in \mathcal{C}(N) \mid PD(l) = p_N + (2g - 2)B_N + \beta_N \Sigma_N \text{ with } \beta_N \in \mathbb{Z} \}.$$

Then we get:

Theorem 3 (Morgan-Szabó-Taubes). In the situation above we have

$$\sum_{k'\in\mathcal{K}(k)}SW_X(k')=\pm\sum SW_M(l_1)SW_N(l_2),$$

where the sum on the right extends over those $(l_1, l_2) \in \mathcal{K}_M(k) \times \mathcal{K}_N(k)$ with $\beta_X = \beta_M + \beta_N + 2$.

Note that if M and N are of simple type there is for a given characteristic class k at most one non-zero summand on the right hand side of the formula; see also [14].

Finally, if M and N are symplectic and the surfaces Σ_M and Σ_N symplectically embedded then the fibre sum X admits a symplectic structure for all gluing diffeomorphisms ϕ as above [5, 12]. There is a formula for the canonical class of X that in general depends on the choice of C. Let X_0 denote the fibre sum with C = 0, corresponding to the gluing diffeomorphism that identifies the push-offs of the surfaces.

Theorem 4. Choose a normal form basis for $H_2(X;\mathbb{Z})$ as in Theorem 1. Suppressing Poincaré duality the canonical class of X is given by

(1)
$$K_X = \overline{K_M} + \overline{K_N} + \sum_{i=1}^c r_i R_i + b_X B_X + \sigma_X \Sigma_X,$$

where

$$\overline{K_M} = K_M - (2g - 2)B_M - (K_M B_M - (2g - 2)B_M^2)\Sigma_M \in P(M)$$

$$\overline{K_N} = K_N - (2g - 2)B_N - (K_N B_N - (2g - 2)B_N^2)\Sigma_N \in P(N)$$

$$r_i = K_X S_i = K_{X_0} S_i - a_i (K_N B_N + 1 - (2g - 2)B_N^2)$$

$$b_X = 2g - 2$$

$$\sigma_X = K_M B_M + K_N B_N + 2 - (2g - 2)(B_M^2 + B_N^2).$$

In evaluating $K_{X_0}S_i$ we choose the basis of rim tori in X_0 determined by the basis in X and a corresponding dual basis of vanishing surfaces.

The integers a_i are defined as follows: Let $\alpha_1, \ldots, \alpha_c$ denote a basis for the kernel of the homomorphism $i_M \oplus i_N$ on $H_1(\Sigma; \mathbb{Z})$. Then

$$a_i = \langle C, \alpha_i \rangle.$$

4. Invariants of the iterated fibre sums M(n)

Let M' denote an algebraic surface with a Lefschetz pencil and $H_1(M'; \mathbb{Z}) = 0$ and $M \to \mathbb{CP}^1$ the algebraic Lefschetz fibration on the blow-up with fibre of genus g. Using two copies M_1 and M_2 of M we see that the homomorphism

$$i_{M_1} \oplus i_{M_2}$$

on $H_1(\Sigma; \mathbb{Z})$ maps to zero, hence

 $H_1(M(2);\mathbb{Z}) = 0$

and by induction

$$H_1(M(n);\mathbb{Z}) = 0$$

for all $n \ge 2$. We can also consider a twisted fibre sum

$$M(m, n, C) = M(m) \#_{\Sigma = \Sigma} M(n)$$

defined by a gluing diffeomorphism that is determined by a cohomology class C in $H^1(\Sigma; \mathbb{Z})$. There is a diffeomorphism

$$M(m, n, 0) \cong M(m+n).$$

The same argument as above shows that

$$H_1(M(m, n, C); \mathbb{Z}) = 0$$

for all $m, n \geq 1$.

More specifically if M' is simply-connected, then the complement of a general fibre in M is also simply-connected, because the meridian to the surface bounds an embedded disk, coming from one of the exceptional spheres. Hence all fibre sums M(n) and M(m, n, C) are simply-connected.

The Euler characteristic and signature of the fibre sum X = M(n) are given by

$$e(X) = ne(M) + (n-1)(4g-4)$$

$$\sigma(X) = n\sigma(M),$$

and similarly for the twisted fibre sum. To describe the second homology of M(2) we choose in M for the surface B_M one of the exceptional spheres, which form sections for the fibration. In X = M(2)these spheres sew together to define a sphere B_X of self-intersection -2. According to the second Lefschetz Hyperplane theorem the vanishing cycles generate the first homology of the general fibre. By our choice of gluing the corresponding vanishing disks pair up in both summands of the fibre sum to define a set of vanishing classes given by embedded (-2)-spheres, called vanishing spheres. With Proposition 2 we get:

Proposition 5. There exists a basis S_1, \ldots, S_{2g} for the group S'(M(2)) of vanishing surfaces which are linear combinations of embedded vanishing spheres and rim tori.

Hence the intersection form of X = M(2) looks like

$$H_2(M(2);\mathbb{Z}) = P(M_1) \oplus P(M_2) \oplus \bigoplus_{i=1}^{2g} \begin{pmatrix} S_i^2 & 1\\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} -2 & 1\\ 1 & 0 \end{pmatrix}.$$

By induction we get

$$H_2(M(n);\mathbb{Z}) = \bigoplus_{i=1}^n P(M_i) \oplus \bigoplus_{i=1}^{2g(n-1)} \begin{pmatrix} S_i^2 & 1\\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} -n & 1\\ 1 & 0 \end{pmatrix}.$$

In the induction step we perturb the fibration on M(n-1) without changing the vanishing cycles such that the vanishing spheres become disjoint from the singular fibres. The terms in the middle of the formula are the vanishing surface and dual rim tori pairs. The class B_X in M(n) is represented by a symplectic sphere of self-intersection -n which is a section of the fibration. Note that the second Betti number has the value expected from the formula for the Euler characteristic. A similar formula holds for the fibre sums M(m, n, C). However, in this case the vanishing surfaces are in general no longer linear combinations of vanishing spheres.

We want to determine the canonical class of the symplectic fibre sum M(n).

Theorem 6. The canonical class of X = M(n) is given by

$$K_X = \sum_{i=1}^n \overline{K_{M_i}} + (2g-2)B_X + ((n-2) + (2g-2)n)\Sigma_X,$$

where

$$\overline{K_{M_i}} = (K_M + \Sigma_M) - (2g - 2)(B_M + \Sigma_M) \in P(M_i)$$

for all i = 1, ..., n.

Proof. The proof is by induction using the formula in Theorem 4. We first show that all rim tori coefficients are zero. This is equivalent to showing that $K_X S_i = 0$ for all vanishing classes S_i . Note that the vanishing classes are linear combinations of vanishing spheres and rim tori. Hence we only have to show that $K_X V = 0$ for all vanishing spheres V and $K_X T = 0$ for all rim tori T. The

sphere V has a dual rim torus R that intersects it in a single positive transverse point. Smoothing the intersection between V and the rim torus R we get an embedded torus of square zero. By the adjunction inequality and since K_X is a Seiberg-Witten basic class we have for tori of square zero $K_X R = 0$ and $K_X (V + R) = 0$, hence $K_X V = 0$. By the same argument $K_X T = 0$.

We first check the case n = 1 of the formula we want to prove. We have:

$$K_X = (K_M + \Sigma_M) - (2g - 2)(B_M + \Sigma_M) + (2g - 2)B_M + (-1 + (2g - 2))\Sigma_M = K_M.$$

Suppose that $n \ge 2$ and the formula is correct for n-1. Write N = M(n-1) and consider the fibre sum $X = M \#_{\Sigma_M = \Sigma_N} N$. The surface B_M is an exceptional sphere in M and B_N is a symplectic sphere of self-intersection -(n-1) from the previous step. Using the adjunction formula we have $K_M B_M = -1$, hence

$$\overline{K_M} = K_M - (2g - 2)B_M - (-1 + (2g - 2))\Sigma_M$$

= $(K_M + \Sigma_M) - (2g - 2)(B_M + \Sigma_M)$
= $\overline{K_{M_n}}$,

and similarly $K_N B_N = n - 3$, hence

$$\overline{K_N} = K_N - (2g - 2)B_N - ((n - 3) + (2g - 2)(n - 1))\Sigma_N$$

= $\sum_{i=1}^{n-1} \overline{K_{M_i}} + (2g - 2)B_N + ((n - 3) + (2g - 2)(n - 1))\Sigma_N$
- $(2g - 2)B_N - ((n - 3) + (2g - 2)(n - 1))\Sigma_N$
= $\sum_{i=1}^{n-1} \overline{K_{M_i}}.$

We also have

$$b_X = 2g - 2$$

$$\sigma_X = -1 + (n - 3) + 2 - (2g - 2)(-1 - (n - 1))$$

$$= (n - 2) + (2g - 2)n.$$

Adding the terms proves the claim.

Remark 7. Note that for g = 1 and M equal to the elliptic surface E(1) with general fibre F we have $K_M + \Sigma_M = -F + F = 0$. Hence we get the well-known formula $K_X = (n-2)F$ for the canonical class of X = E(n).

In the general case we have for the (maximal) divisibility of the canonical class:

Corollary 8. The divisibility of the canonical class K_X of X = M(n) is the greatest common divisor of n - 2 and the divisibility of the class $K_M + \Sigma_M \in H^2(M; \mathbb{Z})$.

Proof. The greatest common divisor of n - 2 and the divisibility of $K_M + \Sigma_M$ divides K_X . This follows because this number also divides $2g - 2 = (K_M + \Sigma_M)\Sigma_M$ by the adjunction formula. The number then divides all terms in the formula in Theorem 6.

Conversely, let δ denote the divisibility of K_X . It is clear that δ divides 2g-2 since $K_X \Sigma_X = 2g-2$ by the adjunction formula. We have

$$K_X B_X = n - 2$$

since B_X is a symplectic sphere of self-intersection -n. This implies that δ divides also n - 2. The integer δ also has to divide every term $\overline{K_{M_i}}$. This shows that it divides the class $K_M + \Sigma_M$, proving the claim.

Remark 9. Since the complex curve Σ_M in the blow-up $M = M' \# r \overline{\mathbb{CP}}^2 \to \mathbb{CP}^1$ is the proper transform of a curve $\Sigma_{M'}$ in M', the divisibility of $K_M + \Sigma_M$ is equal to the divisibility of $K_{M'} + \Sigma_{M'}$. In fact, both classes are equal because the canonical class and the class of the proper transform are given by

$$\Sigma_M = \Sigma_{M'} - E_1 - \ldots - E_r$$
$$K_M = K_{M'} + E_1 + \ldots + E_r,$$

where E_i denotes the exceptional spheres.

We can also determine the canonical class of the fibre sum M(m, n, C).

Theorem 10. The canonical class of X = M(m, n, C) is given by

$$K_X = \sum_{i=1}^{m+n} \overline{K_{M_i}} + \sum_{i=1}^{2g} r_i R_i + (2g-2)B_X + ((m+n-2) + (2g-2)(m+n))\Sigma_X,$$

where

$$\overline{K_{M_i}} = (K_M + \Sigma_M) - (2g - 2)(B_M + \Sigma_M) \in P(M_i)$$

for all $i = 1, \ldots, m + n$ and

$$c_i = -a_i((2g-1)n-1)$$

for the coefficients a_i determined by the class C as above for a basis $\alpha_1, \ldots, \alpha_{2g}$ of $H_1(\Sigma; \mathbb{Z})$.

Proof. The proof follows from the formula in Theorem 4, because $X_0 = M(m+n)$, hence $K_{X_0}S_i = 0$, and B_N is a symplectic sphere of self-intersection -n.

The following is an immediate consequence which we will use later in deriving an obstruction to extending diffeomorphisms:

Corollary 11. The divisibility of the canonical class K_X of X = M(m, n, C) is the greatest common divisor of m + n - 2, a((2g - 1)n - 1) and the class $K_M + \Sigma_M$, where a denotes the divisibility of the class C.

Note that the divisibility of the class C is the greatest common divisor of the integers a_i . We can use this corollary for example to determine when the manifold M(m, n, C) is spin.

Remark 12. One can show that the divisibility is also equal to the greatest common divisor of m + n - 2, a((2g - 1)m - 1) and $K_M + \Sigma_M$, as required by symmetry.

Finally, we determine some of the Seiberg-Witten invariants of the manifolds M(n) in the special case that M' is a minimal surface of general type. Surfaces of general type are algebraic [2]. We have the following lemma.

Lemma 13. Let M' be a smooth minimal surface of general type embedded in some complex projective space and $\Sigma_{M'}$ a transverse hyperplane section. Then the genus of $\Sigma_{M'}$ is at least two. *Proof.* We have $\Sigma_{M'}^2 = r \ge 1$, where r denotes the degree of M'. Since M' is minimal and of general type we have $K_{M'}C \ge 0$ for all smooth curves C with equality if and only if C is a rational (-2)-curve [2]. Hence $K_{M'}\Sigma_{M'} \ge 1$ and the adjunction formula implies that the genus of $\Sigma_{M'}$ is at least two.

For simplicity we also assume in the following that $b_2^+(M') > 1$. First note that the Poincaré dual of a basic class of M(n) has no rim tori component: This follows as before from the adjunction inequality of Seiberg-Witten theory because the vanishing surfaces are linear combinations of vanishing spheres and rim tori; see [4] for a similar argument. Hence the right hand side in Theorem 3 has only one term. It follows from [14] that the intersection of a basic class of M(n) with Σ_X is either equal to $\pm (2g - 2)$ or zero.

Theorem 14. Suppose that M' is a minimal algebraic surface of general type with $H_1(M'; \mathbb{Z}) = 0$ and $b_2^+ > 1$. Then the only Seiberg-Witten basic class up to sign of X = M(n) which has non-zero intersection with the fibre Σ_X is the canonical class K_X .

Proof. We consider the argument for M(2). The general case follows similarly. By the blow-up formula for the Seiberg-Witten invariants [3] the basic classes of $M = M' \# r \overline{\mathbb{CP}}^2$ are given by [16]

$$L = \pm (K_{M'} \pm E_1 \pm \ldots \pm E_r).$$

Note that on the right hand side of the Morgan-Szabó-Taubes formula only the basic classes on M with $L\Sigma_M = 2g - 2$ are relevant. Suppose that *i* exceptional sphere summands in the bracket have negative sign and r - i positive. We have

$$2g - 2 = K_{M'} \Sigma_{M'} + r.$$

Since M' is minimal and of general type and $\Sigma_{M'}$ is a smooth complex curve of non-zero genus we have $K_{M'}\Sigma_{M'} > 0$, which implies $i \leq r < 2g - 2$. We get:

$$|L\Sigma_M| = |K_{M'}\Sigma_{M'} + r - 2i| = |2g - 2 - 2i|.$$

This can be 2g - 2 if and only if i = 0. Hence the relevant basic class of M for the right hand side of the Morgan-Szabó-Taubes formula is $L = K_M$. We then see that the only basic class of M(2) which has intersection 2g - 2 with the fibre is the canonical class.

5. EXTENSION OF DIFFEOMORPHISMS

Suppose that M' is a minimal surface of general type with $H_1(M'; \mathbb{Z}) = 0$ and $b_2^+ > 1$. Let M(n) be a fibre sum as above and Σ a general fibre in M(n) with tubular neighbourhood $\nu\Sigma$. The neighbourhood has a natural framing $\Sigma \times D^2$ given by the fibration. We are interested in orientation preserving self-diffeomorphisms ψ of $\partial \nu\Sigma$ which preserve the circle fibration and cover the identity. As before, the diffeomorphism ψ has up to isotopy the form

$$\psi: \Sigma \times S^1 \longrightarrow \Sigma \times S^1$$
$$(x, \alpha) \mapsto (x, C(x) \cdot \alpha),$$

where $C : \Sigma \to S^1$ is a smooth map. The diffeomorphism is determined up to isotopy by the cohomology class $C^* d\alpha \in H^1(\Sigma; \mathbb{Z})$, denoted by C.

We want to derive an obstruction so that ψ does not extend to an orientation preserving selfdiffeomorphism of the complement $M(n) \setminus int \nu \Sigma$. **Theorem 15.** Let d denote the divisibility of the class $K_{M'} + \Sigma_{M'}$ and a the divisibility of the class C. If d does not divide a(n-1) then ψ does not extend to an orientation preserving self-diffeomorphism of the complement $M(n) \setminus int \nu \Sigma$.

Proof. Suppose that ψ extends to a self-diffeomorphism Ψ of the complement. We can use ψ to form a twisted fibre sum M(m, n, C) for some integer $m \ge 1$. Since ψ extends on the complement there is an orientation preserving diffeomorphism $M(m, n, C) \cong M(m + n)$. Let X' = M(m, n, C) and X = M(m + n). Note that the diffeomorphism is the identity on $M(m)_0$. Hence it maps the surface $\Sigma_{X'}$ in X' given by the standard fibre in the boundary of the tubular neighbourhood in M(m) to the standard fibre Σ_X in M(m + n). On the $M(n)_0$ side in X' the surface $\Sigma_{X'}$ is a twisted copy of the standard fibre and also gets mapped to the fibre Σ_X in X. The canonical class of X' has intersection 2g - 2 with the symplectic surface $\Sigma_{X'}$. It has to map to a Seiberg-Witten basic class of X having the same intersection 2g - 2 with the fibre Σ_X , since the surfaces get identified and the canonical class is basic [15]. There is a unique such basic class, the canonical class of X. It follows that both canonical classes have to match under the diffeomorphism, in particular they must have the same divisibility. This will imply that d divides a(n - 1).

The divisibility of the canonical class of X is the greatest common divisor of

$$m+n-2$$
 and d .

The divisibility of the canonical class of X' is the greatest common divisor of

$$m + n - 2$$
, d and $a((2g - 1)n - 1)$.

If both are the same the greatest common divisor of m + n - 2 and d must divide a((2g - 1)n - 1). We can choose for m any positive integer. In particular, we can arrange that the greatest common divisor of m + n - 2 and d is equal to d. Hence d has to divide a((2g - 1)n - 1). The integer d is the divisibility of $K_{M'} + \Sigma_{M'}$. The adjunction formula shows that d divides 2g - 2. It follows that ddivides a((2g - 1)n - 1) if and only if it divides a(n - 1). This completes the proof.

Remark 16. The corresponding statement is true for elliptic surfaces and shows that if a diffeomorphism extends over $E(n) \setminus \text{int } \nu \Sigma$ then d divides a(n-1). Note that in the case of E(n) the integer d is zero, see Remark 7. This implies that we must have either n = 1, and we are in the case of E(1), or a = 0 and hence C = 0, which is the case of a trivial diffeomorphism that preserves the torus fibration. Hence we get the same obstruction that is known for elliptic surfaces, see Theorem 8.3.11 in [6].

6. CONSTRUCTION OF SOME EXAMPLES

Let M' be a minimal complex surface of general type. Then M' is algebraic and an embedding into some projective space \mathbb{CP}^N is determined by a very ample line bundle E on M'. Under such an embedding, a transverse hyperplane section of M' will be a surface $\Sigma_{M'}$ on M' representing the Poincaré dual of $c_1(E)$. We want to prove the following:

Proposition 17. For each integer $d \ge 1$ there exists an embedding of M' into some projective space \mathbb{CP}^N such that the class $K_{M'} + \Sigma_{M'}$ is divisible by d.

Using this proposition we can construct many Lefschetz fibrations M(n) such that certain diffeomorphisms on the boundary of the tubular neighbourhood of a general fibre do not extend over the complement according to Theorem 15.

We need some preparations: If S is an ample line bundle on M', then kS is very ample for all $k \ge k_0$ for some integer k_0 [11, Theorem 1.2.6]. Let L be an ample line bundle on M'. We want to determine when the line bundle of the form $S = K_{M'} + sL$ for integers $s \ge 1$ is ample.

Lemma 18. Let L be an ample line bundle on M'. Then there exists an integer $s_0 \ge 1$ such that $S = K_{M'} + sL$ is ample for all $s \ge s_0$.

Proof. According to the Nakai-Moishezon criterion [2, 9] the line bundle S is ample if and only if $S^2 > 0$ and SC > 0 for every irreducible curve C on M'. The canonical class of an algebraic surface of general type satisfies $K_{M'}C \ge 0$ for all irreducible curves C, with equality if and only if C is a rational (-2)-curve. Since we assumed that L is ample, we have SC > 0 for all integers $s \ge 1$ and every curve C. Moreover, since L is ample we also have $L^2 > 0$. Hence the second condition $S^2 = K_{M'}^2 + 2sK_{M'}L + s^2L^2 > 0$ will certainly hold if we choose s large enough. In fact, under our assumptions $s \ge 0$ will suffice.

We can now prove Proposition 17.

Proof. Let $S = K_{M'} + sL$ be ample and kS very ample. Then the hyperplane section $\Sigma_{M'}$ under the embedding determined by kS into a projective space \mathbb{CP}^N represents the class $kK_{M'} + ksL$. Let $d \ge 1$ be some integer. We want to choose s and k such that the class $K_{M'} + \Sigma_{M'}$ is divisible by d. First choose s such that $S = K_{M'} + sL$ is ample and d divides s. Then choose k large enough such that kS is very ample and d divides k + 1. Then d also divides the class $K_{M'} + \Sigma_{M'}$.

REFERENCES

- A. Andreotti, T. Frankel, *The second Lefschetz theorem on hyperplane sections*, in: Global Analysis (Papers in Honor of K. Kodaira), 1–20 Univ. Tokyo Press, Tokyo 1969.
- W. P. Barth, K. Hulek, C. A. M. Peters, A. Van de Ven, *Compact complex surfaces*, Second edition, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, Springer-Verlag, Berlin, 2004.
- 3. R. Fintushel, R. J. Stern, *Immersed spheres in 4-manifolds and the immersed Thom conjecture*, Turkish J. Math. **19**, 145–157 (1995).
- 4. R. Fintushel, R. J. Stern, *Families of simply connected 4-manifolds with the same Seiberg-Witten invariants*, Topology **43**, 1449–1467 (2004).
- 5. R. E. Gompf, A new construction of symplectic manifolds, Ann. of Math. 142, 527-595 (1995).
- R. E. Gompf, A. I. Stipsicz, 4-manifolds and Kirby calculus, Graduate Studies in Mathematics, 20. Providence, Rhode Island. American Mathematical Society 1999.
- 7. M. J. D. Hamilton, On symplectic 4-manifolds and contact 5-manifolds, Ph. D. Thesis, Munich 2008, http://edoc.ub.uni-muenchen.de/8779/
- 8. M. J. D. Hamilton, Generalized fibre sums of 4-manifolds and the canonical class, preprint arXiv:0907.2671v3.
- 9. R. Hartshorne, Algebraic Geometry, Graduate Texts in Mathematics 52, Springer-Verlag, New York-Heidelberg, 1977.
- 10. K. Lamotke, The topology of complex projective varieties after S. Lefschetz, Topology 20, 15–51 (1981).
- 11. R. Lazarsfeld, Positivity in algebraic geometry. I. Classical setting: line bundles and linear series, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, Springer-Verlag, Berlin-Heidelberg, 2004.
- 12. J. D. McCarthy, J. G. Wolfson, Symplectic normal connect sum, Topology 33, 729-764 (1994).
- J. Morgan, Z. Szabó, C. H. Taubes, A product formula for the Seiberg-Witten invariants and the generalized Thom conjecture, J. Differential Geom. 44, 706–788 (1996).
- V. Muñoz, B.-L. Wang, Seiberg-Witten-Floer homology of a surface times a circle for non-torsion spin^C structures, Math. Nachr. 278, 844–863 (2005).
- 15. C. H. Taubes, The Seiberg-Witten invariants and symplectic forms, Math. Res. Lett. 1, no. 6, 809-822 (1994).
- 16. E. Witten, Monopoles and four-manifolds, Math. Res. Lett. 1, no. 6, 769-796 (1994).

M. J. D. Hamilton Fachbereich Mathematik Universität Stuttgart Pfaffenwaldring 57 70569 Stuttgart Germany **E-Mail:** mark.hamilton@math.lmu.de

12

Erschienene Preprints ab Nummer 2007/2007-001

Komplette Liste: http://www.mathematik.uni-stuttgart.de/preprints

- 2012-010 Hamilton, M.J.D.: The closure of the symplectic cone of elliptic surfaces
- 2012-009 Hamilton, M.J.D.: Iterated fibre sums of algebraic Lefschetz fibrations
- 2012-008 Hamilton, M.J.D.: The minimal genus problem for elliptic surfaces
- 2012-007 *Ferrario, P:* Partitioning estimation of local variance based on nearest neighbors under censoring
- 2012-006 Stroppel, M.: Buttons, Holes and Loops of String: Lacing the Doily
- 2012-005 Hantsch, F.: Existence of Minimizers in Restricted Hartree-Fock Theory
- 2012-004 Grundhöfer, T.; Stroppel, M.; Van Maldeghem, H.: Unitals admitting all translations
- 2012-003 Hamilton, M.J.D.: Representing homology classes by symplectic surfaces
- 2012-002 *Hamilton, M.J.D.:* On certain exotic 4-manifolds of Akhmedov and Park
- 2012-001 Jentsch, T.: Parallel submanifolds of the real 2-Grassmannian
- 2011-028 Spreer, J.: Combinatorial 3-manifolds with cyclic automorphism group
- 2011-027 *Griesemer, M.; Hantsch, F.; Wellig, D.:* On the Magnetic Pekar Functional and the Existence of Bipolarons
- 2011-026 Müller, S.: Bootstrapping for Bandwidth Selection in Functional Data Regression
- 2011-025 *Felber, T.; Jones, D.; Kohler, M.; Walk, H.:* Weakly universally consistent static forecasting of stationary and ergodic time series via local averaging and least squares estimates
- 2011-024 Jones, D.; Kohler, M.; Walk, H.: Weakly universally consistent forecasting of stationary and ergodic time series
- 2011-023 *Györfi, L.; Walk, H.:* Strongly consistent nonparametric tests of conditional independence
- 2011-022 *Ferrario, P.G.; Walk, H.:* Nonparametric partitioning estimation of residual and local variance based on first and second nearest neighbors
- 2011-021 Eberts, M.; Steinwart, I.: Optimal regression rates for SVMs using Gaussian kernels
- 2011-020 *Frank, R.L.; Geisinger, L.:* Refined Semiclassical Asymptotics for Fractional Powers of the Laplace Operator
- 2011-019 Frank, R.L.; Geisinger, L.: Two-term spectral asymptotics for the Dirichlet Laplacian on a bounded domain
- 2011-018 Hänel, A.; Schulz, C.; Wirth, J.: Embedded eigenvalues for the elastic strip with cracks
- 2011-017 Wirth, J.: Thermo-elasticity for anisotropic media in higher dimensions
- 2011-016 Höllig, K.; Hörner, J.: Programming Multigrid Methods with B-Splines
- 2011-015 Ferrario, P.: Nonparametric Local Averaging Estimation of the Local Variance Function
- 2011-014 *Müller, S.; Dippon, J.:* k-NN Kernel Estimate for Nonparametric Functional Regression in Time Series Analysis
- 2011-013 Knarr, N.; Stroppel, M.: Unitals over composition algebras
- 2011-012 *Knarr, N.; Stroppel, M.:* Baer involutions and polarities in Moufang planes of characteristic two
- 2011-011 Knarr, N.; Stroppel, M.: Polarities and planar collineations of Moufang planes
- 2011-010 Jentsch, T.; Moroianu, A.; Semmelmann, U.: Extrinsic hyperspheres in manifolds with special holonomy
- 2011-009 Wirth, J.: Asymptotic Behaviour of Solutions to Hyperbolic Partial Differential Equations
- 2011-008 Stroppel, M.: Orthogonal polar spaces and unitals
- 2011-007 *Nagl, M.:* Charakterisierung der Symmetrischen Gruppen durch ihre komplexe Gruppenalgebra
- 2011-006 *Solanes, G.; Teufel, E.:* Horo-tightness and total (absolute) curvatures in hyperbolic spaces
- 2011-005 Ginoux, N.; Semmelmann, U.: Imaginary Kählerian Killing spinors I

- 2011-004 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic *D*-Scales: Part II Gain-Scheduled Control
- 2011-003 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic *D*-Scales: Part I Robust Control
- 2011-002 Alexandrov, B.; Semmelmann, U.: Deformations of nearly parallel G₂-structures
- 2011-001 Geisinger, L.; Weidl, T.: Sharp spectral estimates in domains of infinite volume
- 2010-018 Kimmerle, W.; Konovalov, A.: On integral-like units of modular group rings
- 2010-017 *Gauduchon, P.; Moroianu, A.; Semmelmann, U.:* Almost complex structures on quaternion-Kähler manifolds and inner symmetric spaces
- 2010-016 Moroianu, A.; Semmelmann, U.: Clifford structures on Riemannian manifolds
- 2010-015 *Grafarend, E.W.; Kühnel, W.:* A minimal atlas for the rotation group SO(3)
- 2010-014 Weidl, T.: Semiclassical Spectral Bounds and Beyond
- 2010-013 Stroppel, M.: Early explicit examples of non-desarguesian plane geometries
- 2010-012 *Effenberger, F.:* Stacked polytopes and tight triangulations of manifolds
- 2010-011 *Györfi, L.; Walk, H.:* Empirical portfolio selection strategies with proportional transaction costs
- 2010-010 Kohler, M.; Krzyżak, A.; Walk, H.: Estimation of the essential supremum of a regression function
- 2010-009 *Geisinger, L.; Laptev, A.; Weidl, T.:* Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010-008 *Poppitz, S.; Stroppel, M.:* Polarities of Schellhammer Planes
- 2010-007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.:* Non-existence of isomorphisms between certain unitals
- 2010-006 *Höllig, K.; Hörner, J.; Hoffacker, A.:* Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods
- 2010-005 *Kaltenbacher, B.; Walk, H.:* On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010-004 Kühnel, W.; Solanes, G.: Tight surfaces with boundary
- 2010-003 *Kohler, M; Walk, H.:* On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010-002 *Gulde, M.; Stroppel, M.:* Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010-001 *Leitner, F.:* Examples of almost Einstein structures on products and in cohomogeneity one
- 2009-008 Griesemer, M.; Zenk, H.: On the atomic photoeffect in non-relativistic QED
- 2009-007 *Griesemer, M.; Moeller, J.S.:* Bounds on the minimal energy of translation invariant n-polaron systems
- 2009-006 *Demirel, S.; Harrell II, E.M.:* On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009-005 Bächle, A, Kimmerle, W.: Torsion subgroups in integral group rings of finite groups
- 2009-004 Geisinger, L.; Weidl, T.: Universal bounds for traces of the Dirichlet Laplace operator
- 2009-003 Walk, H.: Strong laws of large numbers and nonparametric estimation
- 2009-002 Leitner, F.: The collapsing sphere product of Poincaré-Einstein spaces
- 2009-001 *Brehm, U.; Kühnel, W.:* Lattice triangulations of E^3 and of the 3-torus
- 2008-006 *Kohler, M.; Krzyżak, A.; Walk, H.:* Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps
- 2008-005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.:* Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems

- 2008-004 *Leitner, F.:* Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008-003 Effenberger, F.; Kühnel, W.: Hamiltonian submanifolds of regular polytope
- 2008-002 *Hertweck, M.; Höfert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups PSL(2,q)
- 2008-001 Kovarik, H.; Vugalter, S.; Weidl, T.: Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007-006 Weidl, T.: Improved Berezin-Li-Yau inequalities with a remainder term
- 2007-005 Frank, R.L.; Loss, M.; Weidl, T.: Polya's conjecture in the presence of a constant magnetic field
- 2007-004 *Ekholm, T.; Frank, R.L.; Kovarik, H.:* Eigenvalue estimates for Schrödinger operators on metric trees
- 2007-003 Lesky, P.H.; Racke, R.: Elastic and electro-magnetic waves in infinite waveguides
- 2007-002 Teufel, E.: Spherical transforms and Radon transforms in Moebius geometry
- 2007-001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions