# Universität Stuttgart

# Fachbereich Mathematik

### Convergence of Adaptive Finite Elements for Control Constrained Optimal Control Problems

Kristina Kohls, Arnd Rösch, Kunibert G. Siebert

Preprint 2013/011

Fachbereich Mathematik Fakultät Mathematik und Physik Universität Stuttgart Pfaffenwaldring 57 D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de
WWW: http://www.mathematik.uni-stuttgart.de/preprints

ISSN 1613-8309

C Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors. LaTEX-Style: Winfried Geis, Thomas Merkle

### **Convergence of Adaptive Finite Elements for Control Constrained Optimal Control Problems**

Kristina Kohls, Arnd Rösch and Kunibert G. Siebert

**Abstract.** We summarize our findings in the analysis of adaptive finite element methods for the efficient discretization of control constrained optimal control problems. We particularly focus on convergence of the adaptive method, i. e., we show that the sequence of adaptively generated discrete solutions converges to the true solution. We restrict the presentation to a simple model problem to highlight the key components of the convergence proof and comment on generalizations of the presented result.

**Keywords.** Adaptive finite elements, a posteriori error estimators, convergence analysis, optimal control, control constraints.

#### 1. Statement of the Main Result

In this summary we analyze adaptive finite element discretizations for control constrained optimal control problems of the form

$$\min_{\substack{(u,y)\in\mathbb{U}^{\mathrm{ad}}\times\mathbb{Y}\\ \mathrm{subject to}}} \frac{1}{2} \|y-y_d\|_{\mathbb{U}}^2 + \frac{\alpha}{2} \|u\|_{\mathbb{U}}^2 \tag{1.1}$$
subject to  $y\in\mathbb{Y}$ :  $\mathcal{B}[y,v] = \langle u,v \rangle$   $v\in\mathbb{Y}$ .

In order to highlight the basic ideas of our convergence analysis we focus on the most simple model problem in the following setting. We let  $\Omega \subset \mathbb{R}^d$  be a bounded domain that is meshed exactly by some conforming initial triangulation  $\mathcal{G}_0$ . We consider distributed control in  $\mathbb{U} = L_2(\Omega)$  with a non-empty, convex, and closed subset  $\mathbb{U}^{\mathrm{ad}}$  of admissible controls. We use the  $L_2(\Omega)$  scalar product  $\langle \cdot, \cdot \rangle$  and write  $\| \cdot \|_{\mathbb{U}} = \| \cdot \|_{2;\Omega}$  for its induced norm. The PDE constraint is given by Poisson's problem in the state space  $\mathbb{Y} = \mathring{H}^1(\Omega)$  equipped with norm  $\| \cdot \|_{\mathbb{Y}} = \| \nabla \cdot \|_{2;\Omega}$  and the continuous and coercive bilinear form

$$\mathcal{B}[y, v] = \langle \nabla y, \nabla v \rangle \qquad \forall y, v \in \mathbb{Y}.$$

Finally,  $y_d \in L_2(\Omega)$  is a desired state and  $\alpha > 0$  is some given cost parameter.

Turning to the discretization of (1.1) we denote by  $\mathbb{G}$  the class of all conforming refinements of  $\mathcal{G}_0$  that can be constructed using refinement by bisection [15]. For a given grid  $\mathcal{G} \in \mathbb{G}$  we let  $\mathbb{Y}(\mathcal{G}) \subset \mathbb{Y}$  be a conforming finite element space of piecewise polynomials of fixed degree  $q \in \mathbb{N}$ . We then consider the variational discretization of (1.1) by Hinze [5], i. e., we solve the discretized optimal control problem

 $\min_{\substack{(U,Y)\in\mathbb{U}^{\mathrm{ad}}\times\mathbb{Y}(\mathcal{G})}}\frac{1}{2}\|Y-y_d\|_{\mathbb{U}}^2 + \frac{\alpha}{2}\|U\|_{\mathbb{U}}^2$ subject to  $Y\in\mathbb{Y}(\mathcal{G}): \quad \mathcal{B}[Y,V] = \langle U,V \rangle \quad V\in\mathbb{Y}(\mathcal{G}).$ (1.2)

It is well-known that (1.1) as well as (1.2) admit a unique solution pair  $(\hat{u}, \hat{y})$ , respectively  $(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}})$ ; compare with [11, 17]. Below we additionally utilize the continuous and discrete adjoint states  $\hat{p} \in \mathbb{Y}$ ,  $\hat{P}_{\mathcal{G}} \in \mathbb{Y}(\mathcal{G})$ , and consider the solution triplets  $(\hat{u}, \hat{y}, \hat{p}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y} \times \mathbb{Y}$  and  $(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y}(\mathcal{G}) \times \mathbb{Y}(\mathcal{G})$ .

We use the following adaptive algorithm for approximating the true solution of (1.1). Starting with the initial conforming triangulation  $\mathcal{G}_0$  of  $\Omega$  we execute the standard adaptive loop

SOLVE 
$$\longrightarrow$$
 ESTIMATE  $\longrightarrow$  MARK  $\longrightarrow$  REFINE. (1.3)

In practice, a stopping test is used after ESTIMATE for terminating the iteration; here we shall ignore it for notational convenience.

Assumption 1.1 (Properties of modules). For a given grid  $\mathcal{G} \in \mathbb{G}$  the four used modules have the following properties.

- (1) The output  $(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) := \mathsf{SOLVE}(\mathcal{G}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y}(\mathcal{G}) \times \mathbb{Y}(\mathcal{G})$  is the exact solution of (1.2).
- (2) The output  $\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E)\}_{E \in \mathcal{G}} := \mathsf{ESTIMATE}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); \mathcal{G})$  is a *reliable* and *locally efficient* estimator for the error in the norm  $\|\cdot\|_{\mathbb{U}\times\mathbb{Y}\times\mathbb{Y}}$ . In §2 below we give an example of such an estimator.
- (3) The output  $\mathcal{M} = \mathsf{MARK}(\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E)\}_{E \in \mathcal{G}}, \mathcal{G})$  is a subset of elements subject to refinement. We shall allow any marking strategy such that  $\mathcal{M}$  contains an element holding the maximal indicator, i. e.,

 $\max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) \mid E \in \mathcal{G}\} \le \max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) \mid E \in \mathcal{M}\}.$ 

All practically relevant marking strategies do have this property.

(4) The output  $\mathcal{G}_+ := \mathsf{REFINE}(\mathcal{G}, \mathcal{M}) \in \mathbb{G}$  is a conforming refinement of  $\mathcal{G}$  such that all elements in  $\mathcal{M}$  are bisected at least once, i. e.,  $\mathcal{G}_+ \cap \mathcal{M} = \emptyset$ .

The main contribution of this report is the following convergence result.

**Theorem 1.2 (Main result).** Let  $(\hat{u}, \hat{y}, \hat{p}) \in \mathbb{U}^{ad} \times \mathbb{Y} \times \mathbb{Y}$  be the true solution of (1.1). Suppose that  $\{\hat{U}_k, \hat{Y}_k, \hat{P}_k\}_{k \geq 0} \subset \mathbb{U}^{ad} \times \mathbb{Y} \times \mathbb{Y}$  is any sequence of discrete solutions generated by the adaptive iteration (1.3), where the modules have the properties stated in Assumption 1.1. Then we have

$$\lim_{k \to \infty} \| (\hat{U}_k, \hat{Y}_k, \hat{P}_k) - (\hat{u}, \hat{y}, \hat{p}) \|_{\mathbb{U} \times \mathbb{Y} \times \mathbb{Y}} = 0 \quad and \quad \lim_{k \to \infty} \mathcal{E}_{\mathcal{G}_k} (\hat{U}_k, \hat{Y}_k, \hat{P}_k; \mathcal{G}_k) = 0.$$

The proof of this theorem uses results and ideas from the convergence proofs of Morin, Siebert, and Veeser in [14] and Siebert in [16]. It is a two step procedure presented in §3 and §4. In §3 we utilize basic *stability properties* of the algorithm to show that the sequence of discrete solutions converges to some triplet  $(\hat{u}_{\infty}, \hat{y}_{\infty}, \hat{p}_{\infty})$ . The second step in §4 then relies on the *steering mechanisms* of (1.3), mainly encoded in properties of ESTIMATE and MARK, to finally prove  $(\hat{u}_{\infty}, \hat{y}_{\infty}, \hat{p}_{\infty}) = (\hat{u}, \hat{y}, \hat{p})$ .

We shortly comment on an existing convergence result for constrained optimal control problems given in [2]. It is based on some non-degeneracy assumptions on the continuous and the discrete problems and a smallness assumption on the maximal mesh-size of  $\mathcal{G}_0$ . Our approach does not require any of these assumptions and it is valid for a larger class of adaptive algorithms. In addition, it can easily be extended in several directions; compare with §5.

#### 2. Aposteriori Error Estimation

In this section we shortly summarize our findings from [8, 9] providing a unifying framework for the aposteriori error analysis for control constrained optimal control problems.

First order optimality systems. The analysis in [8] is based on the characterization of the solutions by the first order optimality systems. We let  $S, S^* \colon \mathbb{U} \to \mathbb{Y}$  be the solution operators of the state and the adjoint equation, i.e., for any  $u \in U$  we have

$$Su \in \mathbb{Y}$$
:  $\mathcal{B}[Su, v] = \langle u, v \rangle \quad \forall v \in \mathbb{Y}$  (2.1)

and for any  $g \in \mathbb{U}$  we have

$$S^*g \in \mathbb{Y}: \qquad \mathcal{B}[v, S^*g] = \langle g, v \rangle \qquad \forall v \in \mathbb{Y}.$$
(2.2)

We denote by  $\Pi: \mathbb{U} \to \mathbb{U}^{\mathrm{ad}}$  the nonlinear projection operator such that  $\Pi(p)$  is the best approximation of  $-\frac{1}{\alpha}p$  in  $\mathbb{U}^{\mathrm{ad}}$ , i.e.,

$$\Pi(p) \in \mathbb{U}^{\mathrm{ad}}: \qquad \langle \alpha \Pi(p) + p, \, \Pi(p) - u \rangle \le 0 \qquad \forall u \in \mathbb{U}^{\mathrm{ad}}.$$
(2.3)

Utilizing these operators, the continuous solution  $(\hat{u}, \hat{y}, \hat{p}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y} \times \mathbb{Y}$  is the unique solution of the coupled nonlinear system

$$\hat{y} = S\hat{u}, \qquad \hat{p} = S^*(\hat{y} - y_d), \qquad \hat{u} = \Pi(\hat{p}).$$
 (2.4)

For  $\mathcal{G} \in \mathbb{G}$  we next define the discrete solution operators  $S_{\mathcal{G}}, S_{\mathcal{G}}^* \colon \mathbb{U} \to \mathbb{Y}(\mathcal{G})$ for (2.1) and (2.2), i. e., for any  $u \in \mathbb{U}$  we have

$$S_{\mathcal{G}}u \in \mathbb{Y}(\mathcal{G}): \qquad \mathcal{B}[S_{\mathcal{G}}u, V] = \langle u, V \rangle \qquad \forall V \in \mathbb{Y}(\mathcal{G}),$$
(2.5)

and for any  $g \in \mathbb{U}$  we have

$$S_{\mathcal{G}}^*g \in \mathbb{Y}(\mathcal{G}): \qquad \mathcal{B}[V, S_{\mathcal{G}}^*g] = \langle g, V \rangle \qquad \forall V \in \mathbb{Y}(\mathcal{G}).$$
(2.6)

The discrete solution  $(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y}(\mathcal{G}) \times \mathbb{Y}(\mathcal{G})$  is then uniquely characterized by

$$\hat{Y}_{\mathcal{G}} = S_{\mathcal{G}} \hat{U}_{\mathcal{G}}, \qquad \hat{P}_{\mathcal{G}} = S_{\mathcal{G}}^* (\hat{Y}_{\mathcal{G}} - y_d), \qquad \hat{U}_{\mathcal{G}} = \Pi(\hat{P}_{\mathcal{G}}).$$
(2.7)

Note, that this variational discretization of Hinze requires the evaluation of the *continuous* projection operator  $\Pi$  for discrete functions  $P \in \mathbb{Y}(\mathcal{G})$ .

Coercivity of  $\mathcal{B}$  with constant 1 in combination with the Friedrichs inequality  $||v||_{2;\Omega} \leq C_F ||\nabla v||_{2;\Omega}$  for  $v \in \mathring{H}^1(\Omega)$  yields  $||S||, ||S^*||, ||S_{\mathcal{G}}||, ||S^*_{\mathcal{G}}|| \leq C_F$ .

**Basic error equivalence.** The main obstacle in the aposteriori error analysis encountered for instance in [12, 4] can be explained as follows. One would like to exploit Galerkin orthogonality in the linear state equation (2.1) and the adjoint equation (2.2). However, we observe that triplet  $(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}})$  is the Galerkin approximation to the triplet  $(\hat{u}, \hat{y}, \hat{p})$  but  $\hat{Y}_{\mathcal{G}}$  is not the Galerkin approximation to the solution  $\hat{u}$  of the linear problem (2.1) since we have  $\hat{y} = S\hat{u}$  but not  $\hat{y} = S\hat{U}_{\mathcal{G}}$ . The same argument applies to the adjoint states. This observation shows that we cannot directly employ Galerkin orthogonality for single components of (2.4) and the nonlinearity in (2.3) prevents us from making use of Galerkin orthogonality for the system (2.4). The resort to this problem is given by the following result from [8, Theorem 2.2].

**Proposition 2.1 (Basic error equivalence).** If we set  $\mathbb{W} = \mathbb{U} \times \mathbb{Y} \times \mathbb{Y}$  we have for  $\bar{y} = S\hat{U}_{\mathcal{G}}$  and  $\bar{p} = S^*(\hat{Y}_{\mathcal{G}} - y_d)$  the basic error equivalence

$$\|(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\hat{u}, \hat{p}, \hat{y})\|_{\mathbb{W}} \simeq \|(\hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\bar{y}, \bar{p})\|_{\mathbb{Y} \times \mathbb{Y}}.$$

For the problem under consideration, the constant hidden in  $\simeq$  depends on  $\alpha^{-1}$ . For general  $\mathcal{B}$  it will in addition depend on the inf-sup constant of  $\mathcal{B}$ . Employing this error equivalence it is sufficient to construct a reliable and efficient estimator for the right hand side  $\|(\hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\bar{y}, \bar{p})\|_{\mathbb{Y}\times\mathbb{Y}}$ . The functions  $\bar{y}$  and  $\bar{p}$  are solutions to the *linear* problems (2.1) and (2.2) with given source  $\hat{U}_{\mathcal{G}}$  and  $\hat{Y}_{\mathcal{G}} - y_d$ , respectively. They play a similar role as the *elliptic reconstruction* used in the aposteriori error analysis of parabolic problems; compare with [13].

**Aposteriori error estimation.** We realize that  $\hat{Y}_{\mathcal{G}}$  is the Galerkin approximation to  $\bar{y}$  and  $\hat{P}_{\mathcal{G}}$  the one to  $\bar{p}$ . We therefore can directly employ (existing) estimators for the linear problems (2.1) and (2.2) and their sum then constitutes an estimator for the optimal control problem; compare with [8, Theorem 3.2]. For ease of presentation we focus here on the residual estimator. For any subset  $\mathcal{G}' \subset \mathcal{G}$  we set  $\Omega(\mathcal{G}') := \bigcup_{E \in \mathcal{G}'} E$  and for given  $E \in \mathcal{G}$  we denote by  $\mathcal{N}_{\mathcal{G}}(E) \subset \mathcal{G}$  the subset consisting of E and its direct neighbors. Finally, we indicate by  $\|\cdot\|_{\mathbb{W}(\omega)}$  the natural restriction of  $\|\cdot\|_{\mathbb{W}}$  to a subset  $\omega \subset \Omega$ . We then have the following result.

**Theorem 2.2 (Aposteriori error control).** For  $E \in \mathcal{G}$  we define the indicator

$$\begin{aligned} \mathcal{E}_{\mathcal{G}}^{2}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) &:= h_{E}^{2} \|\Delta \hat{Y}_{\mathcal{G}} + \hat{U}_{\mathcal{G}}\|_{2;E}^{2} + h_{E} \| [\![\nabla \hat{Y}_{\mathcal{G}}]\!]\|_{2;\partial E \cap \Omega}^{2} \\ &+ h_{E}^{2} \|\Delta \hat{P}_{\mathcal{G}} + (\hat{Y}_{\mathcal{G}} - y_{d})\|_{2;E}^{2} + h_{E} \| [\![\nabla \hat{P}_{\mathcal{G}}]\!]\|_{2;\partial E \cap \Omega}^{2} \end{aligned}$$

Then we have the global upper bound

$$\|(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\hat{u}, \hat{p}, \hat{y})\|_{\mathbb{W}}^2 \lesssim \mathcal{E}_{\mathcal{G}}^2((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); \mathcal{G}) := \sum_{E \in \mathcal{G}} \mathcal{E}_{\mathcal{G}}^2((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E).$$

For any  $E \in \mathcal{G}$  we have the local lower bound

$$\mathcal{E}_{\mathcal{G}}^{2}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) \lesssim \|(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\hat{u}, \hat{p}, \hat{y})\|_{\mathbb{W}(\Omega(\mathcal{N}_{\mathcal{G}}(E)))}^{2} + \operatorname{osc}_{\mathcal{G}}^{2}(\hat{U}_{\mathcal{G}}, y_{d}; \mathcal{N}_{\mathcal{G}}(E)),$$

where

$$\operatorname{osc}_{\mathcal{G}}^{2}(\hat{U}_{\mathcal{G}}, y_{d}; E) := h_{E}^{2} \left( \|\hat{U}_{\mathcal{G}} - \mathbb{P}_{\mathcal{G}}\hat{U}_{\mathcal{G}}\|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^{2} + \|y_{d} - \mathbb{P}_{\mathcal{G}}y_{d}\|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^{2} \right)$$

is the typical oscillation term with the  $L_2$ -projection  $\mathbb{P}_{\mathcal{G}}$  onto the set of discontinuous, piecewise polynomials of degree q over  $\mathcal{G}$ .

**Bounds for the residuals.** We shortly comment on the derivation of the estimators for the linear problems and thereby recording an important intermediate estimate. For given  $u \in \mathbb{U}$  we set y = Su and let  $Y = S_{\mathcal{G}}u$  be its Galerkin-approximation in  $\mathbb{Y}(\mathcal{G})$ . Defining the *residual* of the state equation (2.1) by

$$\langle \mathcal{R}(S_{\mathcal{G}}u;u), v \rangle = \langle \mathcal{R}(Y;u), v \rangle := \mathcal{B}[Y,v] - \langle u,v \rangle = \mathcal{B}[Y-y,v] \qquad \forall v \in \mathbb{Y},$$

we find  $\|\mathcal{R}(Y;u)\|_{\mathbb{Y}^*} \simeq \|Y-y\|_{\mathbb{Y}} = \|(S_{\mathcal{G}}-S)u\|_{\mathbb{Y}}.$ 

Employing Galerkin-orthogonality  $\langle \mathcal{R}(Y; u), V \rangle = 0$  for all  $V \in \mathbb{Y}(\mathcal{G})$  and using piecewise integration by parts we deduce for any  $v \in \mathbb{Y}$  and  $V \in \mathbb{V}(\mathcal{G})$  the bound

$$|\langle \mathcal{R}(Y;u), v \rangle| \le \sum_{E \in \mathcal{G}} \|\Delta Y + u\|_{2;E} \|v - V\|_{2;E} + \frac{1}{2} \| [\![\nabla Y]\!]\|_{2;\partial E \cap \Omega} \|v - V\|_{2;\partial E}.$$

Using for  $v \in \mathbb{Y}$  the Scott-Zhang interpolant  $V \in \mathbb{Y}(\mathcal{G})$  one obtains from interpolation estimates in  $H^1$  by standard arguments the upper bound

$$\|Y - y\|_{\mathbb{Y}} \simeq \|\mathcal{R}(Y; u)\|_{\mathbb{Y}^*} \lesssim \left(\sum_{E \in \mathcal{G}} h_E^2 \|\Delta Y + u\|_{2;E}^2 + h_E \|[\![\nabla Y]\!]\|_{2;\partial E \cap \Omega}^2\right)^{1/2}.$$

If v is smooth, i.e.,  $v \in H^2(\Omega) \cap \mathbb{Y}$ , we may employ interpolation estimates in  $H^2$  to obtain the *improved bound* 

$$|\langle \mathcal{R}(Y;u), v \rangle| \lesssim \left( \sum_{E \in \mathcal{G}} h_E^2 (h_E^2 \| \Delta Y + u \|_{2;E}^2 + h_E \| [\![\nabla Y]\!] \|_{2;\partial E \cap \Omega}^2 \right) \right)^{1/2} |v|_{H^2(\Omega)}.$$
(2.8)

Similar arguments apply to the adjoint problem. For given  $g \in \mathbb{U}$  we set  $p = S^*_{\mathcal{G}}g$ and let  $P = S^*_{\mathcal{G}}g$  be its Galerkin-approximation in  $\mathbb{Y}(\mathcal{G})$ . For the residual of (2.2), defined by

$$\langle \mathcal{R}^*(S^*_{\mathcal{G}}g;g), v \rangle = \langle \mathcal{R}^*(P;g), v \rangle := \mathcal{B}[v, P] - \langle g, v \rangle = \mathcal{B}[v, P-p] \qquad \forall v \in \mathbb{Y},$$

5

we have

$$|\langle \mathcal{R}^{*}(P;g), v \rangle| \lesssim \left( \sum_{E \in \mathcal{G}} h_{E}^{2s} \left( h_{E}^{2} \| \Delta P + g \|_{2;E}^{2} + h_{E} \| [\![\nabla P]\!]\|_{2;\partial E \cap \Omega}^{2} \right) \right)^{1/2} |v|_{H^{s+1}(\Omega)}$$
(2.9)

for any  $v \in H^{s+1}(\Omega) \cap \mathbb{Y}$ , s = 0, 1. With s = 0 we may deduce the upper bound for  $\|(S_{\mathcal{G}}^* - S^*)g\|_{\mathbb{Y}} = \|P - p\|_{\mathbb{Y}} \simeq \|\mathcal{R}^*(P;g)\|_{\mathbb{Y}^*}$  The choice s = 1 yields the improved estimate for the adjoint problem. Eqns. (2.8) and (2.9) will become important in §4 to access *local density* of adaptively generated finite element spaces; compare also with [16, Remark 3.4].

#### 3. Convergence 1: Trusting Stability

In this section we start with the convergence analysis, where we first focus on stability properties of the algorithm that do not depend on the particular decisions taken in MARK. Hereafter,  $\{\mathcal{G}_k, (\hat{U}_k, \hat{Y}_k, \hat{P}_k)\}_{k\geq 0}$  is the sequence of grids and discrete solutions generated by (1.3). For ease of notation we use for  $k \geq 0$  the short hands  $\mathbb{Y}_k = \mathbb{Y}(\mathcal{G}_k), \hat{U}_k = \hat{U}_{\mathcal{G}_k}, S_k = S_{\mathcal{G}_k}$  etc.

A first limit. Using piecewise polynomials in combination with refinement by bisection leads to nested spaces, i. e.,  $\mathbb{Y}_k \subset \mathbb{Y}_{k+1}$ . This allows us to define the limiting space

$$\mathbb{Y}_{\infty} = \overline{\bigcup_{k \ge 0} \mathbb{Y}_k}^{\|\cdot\|_{\mathbb{Y}}},$$

which is exactly the space that is approximated by the adaptive iteration. It is closed in  $\mathbb{Y}$  and therefore a Hilbert space. Consequently, the limiting optimal control problem

$$\min_{\substack{(u,y)\in\mathbb{U}^{\mathrm{ad}}\times\mathbb{Y}_{\infty}}}\frac{1}{2}\|y-y_d\|_{\mathbb{U}}^2 + \frac{\alpha}{2}\|u\|_{\mathbb{U}}^2$$
subject to  $y\in\mathbb{Y}_{\infty}: \quad \mathcal{B}[y,v] = \langle u,v\rangle \quad v\in\mathbb{Y}_{\infty}$ 

$$(3.1)$$

admits a unique solution  $(\hat{u}_{\infty}, \hat{y}_{\infty}) \in \mathbb{U}^{\mathrm{ad}} \times \mathbb{Y}_{\infty}$ . If  $S_{\infty}, S_{\infty}^* : \mathbb{U} \to \mathbb{Y}_{\infty}$  denote the solution operators of the state respectively the adjoint equation in  $\mathbb{Y}_{\infty}$  the associated first order optimality system reads

$$\hat{y}_{\infty} = S_{\infty}\hat{u}_{\infty}, \qquad \hat{p}_{\infty} = S_{\infty}^*(\hat{y}_{\infty} - y_d), \qquad \hat{u}_{\infty} = \Pi(\hat{p}_{\infty}). \tag{3.2}$$

We next show that in fact (3.1) is the limiting problem of the adaptive iteration (1.3) in that  $(\hat{U}_k, \hat{Y}_k, \hat{P}_k) \rightarrow (\hat{u}_\infty, \hat{y}_\infty, \hat{p}_\infty)$ . An important ingredient for this proof is the following crucial property of the adaptive algorithm shown in [1, Lemma 6.1] and [14, Lemma 4.2].

**Proposition 3.1 (Convergence of solution operators).** For any  $u, g \in \mathbb{U}$  we have  $S_k u \to S_{\infty} u$  and  $S_k^* g \to S_{\infty}^* g$  in  $\mathbb{Y}$  as  $k \to \infty$ .

We next show convergence  $\hat{U}_k \to \hat{u}_\infty$ . In this step we have to deal with the nonlinearity of the constrained optimal control problem.

**Lemma 3.2 (Convergence of the controls).** The discrete controls  $\{\hat{U}_k\}_{k\geq 0}$  converge strongly to  $\hat{u}_{\infty}$ , *i. e.*,

$$\lim_{k \to \infty} \|\hat{U}_k - \hat{u}_\infty\|_{\mathbb{U}} = 0.$$

*Proof.* Since both  $\hat{U}_k = \Pi(\hat{P}_k)$  and  $\hat{u}_{\infty} = \Pi(\hat{p}_{\infty})$  are feasible, i.e.,  $\hat{U}_k, \hat{u}_{\infty} \in \mathbb{U}^{\mathrm{ad}}$ , the definition of  $\Pi$  in (2.3) yields

$$\begin{aligned} \alpha \|\hat{U}_k - \hat{u}_\infty\|_{2;\Omega}^2 &= \langle \alpha \hat{u}_\infty + \hat{p}_\infty, \, \hat{u}_\infty - \hat{U}_k \rangle + \langle \alpha \hat{U}_k + \hat{P}_k, \, \hat{U}_k - \hat{u}_\infty \rangle \\ &+ \langle \hat{P}_k - \hat{p}_\infty, \, \hat{u}_\infty - \hat{U}_k \rangle \\ &\leq \langle \hat{P}_k - \hat{p}_\infty, \, \hat{u}_\infty - \hat{U}_k \rangle \\ &= \langle S_k^*(\hat{y}_\infty - y_d) - \hat{p}_\infty, \, \hat{u}_\infty - \hat{U}_k \rangle + \langle \hat{P}_k - S_k^*(\hat{y}_\infty - y_d), \, \hat{u}_\infty - \hat{U}_k \rangle. \end{aligned}$$

We next estimate the last two terms separately. For the first one we immediately obtain from  $\hat{p}_{\infty} = S_{\infty}(\hat{y}_{\infty} - y_d)$  by the Cauchy-Schwarz and Young inequalities

$$\begin{aligned} \langle S_k^*(\hat{y}_{\infty} - y_d) - \hat{p}_{\infty}, \, \hat{u}_{\infty} - \hat{U}_k \rangle &= \langle (S_k^* - S_{\infty}^*)(\hat{y}_{\infty} - y_d), \, \hat{u}_{\infty} - \hat{U}_k \rangle \\ &\leq \frac{\alpha}{2} \|\hat{u}_{\infty} - \hat{U}_k\|_{2;\Omega}^2 + \frac{1}{2\alpha} \|(S_k^* - S_{\infty}^*)(\hat{y}_{\infty} - y_d)\|_{2;\Omega}^2 \end{aligned}$$

We next turn to the second term. Employing the definition of the solution operators  $S_k$  and  $S_k^*$  in (2.5) and (2.6) we estimate with  $\hat{P}_k = S_k^*(\hat{Y}_k - y_d) \in \mathbb{Y}_k$  and  $\hat{y}_{\infty} = S_{\infty}\hat{u}_{\infty}$ 

$$\begin{split} \langle \hat{P}_{k} - S_{k}^{*}(\hat{y}_{\infty} - y_{d}), \, \hat{u}_{\infty} - \hat{U}_{k} \rangle &= \langle \hat{u}_{\infty} - \hat{U}_{k}, \, S_{k}^{*}(\hat{Y}_{k} - \hat{y}_{\infty}) \rangle \\ &= \mathcal{B}[S_{k}(\hat{u}_{\infty} - \hat{U}_{k}), \, S_{k}^{*}(\hat{Y}_{k} - \hat{y}_{\infty})] = \langle \hat{Y}_{k} - \hat{y}_{\infty}, \, S_{k}(\hat{u}_{\infty} - \hat{U}_{k}) \rangle \\ &= \langle \hat{Y}_{k} - \hat{y}_{\infty}, \, \hat{y}_{\infty} - \hat{Y}_{k} \rangle + \langle \hat{Y}_{k} - \hat{y}_{\infty}, \, (S_{k} - S_{\infty})\hat{u}_{\infty} \rangle \\ &= - \|\hat{Y}_{k} - \hat{y}_{\infty}\|_{2;\Omega}^{2} + \frac{1}{2} \|\hat{Y}_{k} - \hat{y}_{\infty}\|_{2;\Omega}^{2} + \frac{1}{2} \|(S_{k} - S_{\infty})\hat{u}_{\infty}\|_{2;\Omega}^{2} \\ &\leq \frac{1}{2} \|(S_{k} - S_{\infty})\hat{u}_{\infty}\|_{2;\Omega}^{2}. \end{split}$$

Combining the estimates we have shown

$$\alpha \|\hat{U}_k - \hat{u}_\infty\|_{2;\Omega}^2 \leq \frac{1}{\alpha} \|(S_k^* - S_\infty^*)(\hat{y}_\infty - y_d)\|_{2;\Omega}^2 + \|(S_k - S_\infty)\hat{u}_\infty\|_{2;\Omega}^2 \to 0$$
  
as  $k \to \infty$  by Proposition 3.1. This finishes the proof.  $\Box$ 

Convergence  $(\hat{U}_k, \hat{Y}_k, \hat{P}_k) \to (\hat{u}_\infty, \hat{y}_\infty, \hat{p}_\infty)$  is now a direct consequence of the linear theory in Proposition 3.1.

**Proposition 3.3 (Convergence of discrete solutions).** The Galerkin approximations  $\{(\hat{U}_k, \hat{Y}_k, \hat{P}_k)\}_{k\geq 0}$  converge strongly to the solution  $(\hat{u}_{\infty}, \hat{y}_{\infty}, \hat{p}_{\infty})$  of (3.1), i. e.,

$$\lim_{k \to \infty} \| (\hat{U}_k, \hat{Y}_k, \hat{P}_k) - (\hat{u}_\infty, \hat{y}_\infty, \hat{p}_\infty) \|_{\mathbb{U} \times \mathbb{Y} \times \mathbb{Y}} = 0$$

*Proof.* We already know  $\|\hat{U}_k - \hat{u}_{\infty}\|_{\mathbb{U}} \to 0$  from Lemma 3.2. In combination with Proposition 3.1 this yields for the discrete states

$$\begin{aligned} \|\hat{Y}_{k} - \hat{y}_{\infty}\|_{\mathbb{Y}} &= \|S_{k}\hat{U}_{k} - S_{\infty}\hat{u}_{\infty}\|_{\mathbb{Y}} \leq \|S_{k}(\hat{U}_{k} - \hat{u}_{\infty})\|_{\mathbb{Y}} + \|(S_{k} - S_{\infty})\hat{u}_{\infty}\|_{\mathbb{Y}} \\ &\leq \|S_{k}\| \|\hat{U}_{k} - \hat{u}_{\infty}\|_{\mathbb{U}} + \|(S_{k} - S_{\infty})\hat{u}_{\infty}\|_{\mathbb{Y}} \to 0, \end{aligned}$$

since  $||S_k|| \leq C_F$ . Writing  $\hat{P}_k - \hat{p}_{\infty} = S_k^* (\hat{Y}_k - \hat{y}_{\infty}) + (S_k^* - S_{\infty})(\hat{y}_{\infty} - y_d)$  we finally deduce with the same arguments  $||\hat{P}_k - \hat{p}_{\infty}||_{\mathbb{Y}} \to 0$ .

The convergence of the discrete solutions directly yields a uniform bound on the estimators. The proof follows the ideas in [16, Lemma 3.3] accounting for the situation at hand and using the following important property. Let  $\mathcal{G} \in \mathbb{G}$  be given. The finite overlap of the patches  $\#\mathcal{N}_{\mathcal{G}}(E) \leq 1$  allows us to deduce for any  $g \in L_2(\Omega)$  the bound

$$\sum_{E \in \mathcal{G}} \|g\|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^2 = \sum_{E \in \mathcal{G}} \sum_{E' \in \mathcal{N}_{\mathcal{G}}(E)} \|g\|_{2;E'}^2 \lesssim \sum_{E \in \mathcal{G}} \|g\|_{2;E}^2 = \|g\|_{2;\Omega}^2.$$
(3.3)

The constant solely depends on shape-regularity of  $\mathcal{G}$  and thus on  $\mathcal{G}_0$ .

Lemma 3.4 (Uniform estimator bound). For all  $k \ge 0$  we have

$$\mathcal{E}_k((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); \mathcal{G}_k) \lesssim 1.$$

Proof. A scaled trace inequality in combination with an inverse estimate yields for the error indicators related to the state equation

$$h_E^2 \|\Delta \hat{Y}_k + \hat{U}_k\|_{2;E}^2 + h_E \| [\![\nabla \hat{Y}_k]\!]\|_{2;\partial E_k \cap \Omega}^2 \lesssim \|\nabla \hat{Y}_k\|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^2 + \|\hat{U}_k\|_{2;E}^2.$$

This in turn implies by (3.3)

$$\sum_{E \in \mathcal{G}_k} h_E^2 \|\Delta \hat{Y}_k + \hat{U}_k\|_{2;E}^2 + h_E \| [\![\nabla \hat{Y}_k]\!]\|_{2;\partial E \cap \Omega}^2 \lesssim \|\nabla \hat{Y}_k\|_{2;\Omega}^2 + \|\hat{U}_k\|_{2;\Omega}^2 \lesssim 1,$$

since  $\{\hat{U}_k, \hat{Y}_k\}_{k\geq 0}$  is bounded in  $L_2(\Omega) \times \mathring{H}^1(\Omega)$ . Similar arguments apply to the estimator contribution related to the adjoint problem.

**A second limit.** We next turn to the limit of the piecewise constant mesh-size function  $h_k: \Omega \to \mathbb{R}$  of  $\mathcal{G}_k$  defined by  $h_{k|E} = |E|^{1/d}$ ,  $E \in \mathcal{G}$ . The behavior of the mesh-size function is directly related to the decomposition

$$\mathcal{G}_k^+ := \bigcap_{\ell \ge k} \mathcal{G}_\ell = \{ E \in \mathcal{G}_k \mid E \in \mathcal{G}_\ell \ \forall \ell \ge k \}, \quad \text{and} \quad \mathcal{G}_k^0 := \mathcal{G}_k \setminus \mathcal{G}_k^+.$$

The set  $\mathcal{G}_k^+$  contains all elements that are not refined after iteration k and we observe that the sequence  $\{\mathcal{G}_k^+\}_{k\geq 0}$  is nested, i.e.,  $\mathcal{G}_\ell^+ \subset \mathcal{G}_k^+$  for all  $k \geq \ell$ . The set  $\mathcal{G}_k^0$  contains all elements that are refined at least once more after iteration k; in particular,  $\mathcal{M}_k \subset \mathcal{G}_k^0$ . Decomposing  $\overline{\Omega} = \Omega_k^+ \cup \Omega_k^0 := \Omega(\mathcal{G}_k^+) \cup \Omega(\mathcal{G}_k^0)$  we have the following connection to the behavior of the mesh-size function shown in [14, Lemma 4.3 and Corollary 4.1].

9

Lemma 3.5 (Convergence of the mesh-size functions). The mesh-size functions  $h_k$  converge uniformly to 0 in  $\Omega_k^0$  in the following sense

$$\lim_{k \to \infty} \|h_k \chi_k^0\|_{\infty;\Omega} = \lim_{k \to \infty} \|h_k\|_{\infty;\Omega_k^0} = 0,$$

where  $\chi_k^0 \in L_\infty(\Omega)$  the characteristic function of  $\Omega_k^0$ .

Combining convergence of the discrete solutions with the convergence of the mesh-size functions we see that the adaptive algorithm can monitor progress in the following sense.

Lemma 3.6 (Indicators of marked elements). All indicators of marked elements vanish in the limit, this is,

$$\lim_{k \to \infty} \max \{ \mathcal{E}_{\mathcal{G}}((\dot{U}_{\mathcal{G}}, \dot{Y}_{\mathcal{G}}, \dot{P}_{\mathcal{G}}); E) \mid E \in \mathcal{M}_k \} = 0.$$

Proof. For  $k \ge 0$  we pick up  $E_k \in \arg \max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) \mid E \in \mathcal{M}_k\} \neq \emptyset$ . We follow [16, Lemma 3.4] and show  $\mathcal{E}_k((\hat{U}_k, \hat{Y}_k, \hat{P}_k); E_k) \to 0$ .

Arguing as in the proof to Lemma 3.4 we find for the indicator contribution of the state equation

$$h_E \|\Delta \hat{Y}_k + \hat{U}_k\|_{2;E_k} + h_E^{1/2} \| [\![\nabla \hat{Y}_k]\!]\|_{2;\partial E_k \cap \Omega} \lesssim \|\nabla \hat{Y}_k\|_{2;\Omega(\mathcal{N}_k(E_k))} + \|\hat{U}_k\|_{2;E_k} \\ \le \|\nabla \hat{y}_\infty\|_{2;\Omega(\mathcal{N}_k(E_k))} + \|\hat{u}_\infty\|_{2;E_k} + \|\nabla (\hat{Y}_k - \hat{y}_\infty)\|_{2;\Omega} + \|\hat{U}_k - \hat{u}_\infty\|_{2;\Omega} \to 0$$

as  $k \to \infty$  for the following reasons: By Assumption 1.1 (4) all elements in  $\mathcal{M}_k$  are refined, which implies  $E_k \in \mathcal{G}_k^0$ . Local quasi-uniformity of  $\mathcal{G}_k$  in combination with Lemma 3.5 therefore yields  $|\Omega(\mathcal{N}_k(E_k))| \leq |E_k| \leq ||h_k||_{\infty;\Omega_k^0}^d \to 0$ . Consequently, the first two terms of the right hand side vanish by continuity of  $||\cdot||_{2;\Omega}$  with respect to the Lebesgue measure. The last two terms converge to 0 by Proposition 3.3. The same arguments apply to the indicator contribution of the adjoint equation, which in summary yields  $\mathcal{E}_{\mathcal{G}}((\hat{U}_k, \hat{Y}_k, \hat{P}_k); E_k) \to 0$  as  $k \to \infty$ .

#### 4. Convergence 2: Making the Right Decisions

In this section we verify the main result by showing  $(\hat{U}_k, \hat{Y}_k, \hat{P}_k) \to (\hat{u}, \hat{y}, \hat{p})$  and  $\mathcal{E}_k(\hat{U}_k, \hat{Y}_k, \hat{P}_k; \mathcal{G}_k) \to 0$ . Error convergence requires appropriate decisions in the adaptive iteration, which we have summarized in Assumption 1.1. Estimator convergence is then a consequence of local efficiency as stated in Theorem 2.2.

**Convergence of the indicators.** We first show that the maximal indicator of all elements vanishes in the limit.

Lemma 4.1 (Convergence of the indicators). The maximal indicator vanishes in the limit, this is,

$$\lim_{k \to \infty} \max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}); E) \mid E \in \mathcal{G}_k\} = 0.$$

*Proof.* Combining the assumption on marking in Assumption 1.1 (3) with the behavior of the indicators on marked elements, which we have analyzed in Lemma 3.6, we find

$$\max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_k, \hat{Y}_k, \hat{P}_k); E) \mid E \in \mathcal{G}_k\} \le \max\{\mathcal{E}_{\mathcal{G}}((\hat{U}_k, \hat{Y}_k, \hat{P}_k); E) \mid E \in \mathcal{M}_k\} \to 0$$
  
as  $k \to \infty$ .

**Convergence of the residuals.** We next show that residuals of state and adjoint equation in the limiting first order optimality system (3.2) vanish. The proof adapts the techniques from [16, Proposition 3.1] to the situation at hand.

**Proposition 4.2 (Convergence of the residual).** For the residuals  $\mathcal{R}$  of (2.1) and  $\mathcal{R}^*$  of (2.2) We have

$$\mathcal{R}(\hat{y}_{\infty}; \hat{u}_{\infty}) = \mathcal{R}^*(\hat{p}_{\infty}; \hat{y}_{\infty} - y_d) = 0 \qquad in \ \mathbb{Y}^* = H^{-1}(\Omega).$$

Particularly,  $\hat{y}_{\infty} = S\hat{u}_{\infty}$  and  $\hat{p}_{\infty} = S^*(\hat{y}_{\infty} - y_d)$ .

*Proof.* We prove the claim for  $\mathcal{R}$ . The assertion for  $\mathcal{R}^*$  follows along the same lines. Using a density argument we only have to show  $\langle \mathcal{R}(\hat{y}_{\infty}; \hat{u}_{\infty}), v \rangle = 0$  for all  $v \in H^2(\Omega) \cap \mathring{H}^1(\Omega)$ .

Suppose any pair  $k \ge \ell$ . Then we have the inclusion  $\mathcal{G}_{\ell}^+ \subset \mathcal{G}_k^+ \subset \mathcal{G}_k$  and the sub-triangulation  $\mathcal{G}_k \setminus \mathcal{G}_{\ell}^+$  of  $\mathcal{G}_k$  covers the sub-domain  $\Omega_{\ell}^0 = \Omega(\mathcal{G}_{\ell}^0)$ , i.e., we can write  $\Omega_{\ell}^0 = \Omega(\mathcal{G}_k \setminus \mathcal{G}_{\ell}^+)$ . Moreover,  $\|h_k\|_{\infty;\Omega_{\ell}^+} \lesssim 1$  and  $\|h_k\|_{\infty;\Omega_{\ell}^0} \le \|h_{\ell}\|_{\infty;\Omega_{\ell}^0}$ .

Pick up any  $v \in H^2(\Omega) \cap \mathring{H}^1(\Omega)$  with  $|v|_{H^2(\Omega)} = 1$ . We next utilize the improved bound (2.8) for  $\mathcal{R}$ , decompose  $\mathcal{G}_k = \mathcal{G}_\ell^+ \cup (\mathcal{G}_k \setminus \mathcal{G}_\ell^+)$ , and recall Lemma 3.4 to bound

$$\begin{aligned} \langle \mathcal{R}(\hat{Y}_{k};\hat{U}_{k}), v \rangle^{2} &\lesssim \sum_{E \in \mathcal{G}_{\ell}^{+}} h_{E}^{2} \big( h_{E}^{2} \| \Delta \hat{Y}_{k} + \hat{U}_{k} \|_{2;E}^{2} + h_{E} \| \llbracket \nabla \hat{Y}_{k} \rrbracket \|_{2;\partial E \cap \Omega}^{2} \big) \\ &+ \sum_{E \in \mathcal{G}_{k} \setminus \mathcal{G}_{\ell}^{+}} h_{E}^{2} \big( h_{E}^{2} \| \Delta \hat{Y}_{k} + \hat{U}_{k} \|_{2;E}^{2} + h_{E} \| \llbracket \nabla \hat{Y}_{k} \rrbracket \|_{2;\partial E \cap \Omega}^{2} \big) \\ &\lesssim \mathcal{E}_{k}^{2} ((\hat{U}_{k}, \hat{Y}_{k}, \hat{P}_{k}); \mathcal{G}_{\ell}^{+}) + \| h_{\ell} \|_{\infty;\Omega_{\ell}^{0}}^{2} \mathcal{E}_{k}^{2} ((\hat{U}_{k}, \hat{Y}_{k}, \hat{P}_{k}); \mathcal{G}_{k} \setminus \mathcal{G}_{\ell}^{+}) \\ &\lesssim \mathcal{E}_{k}^{2} ((\hat{U}_{k}, \hat{Y}_{k}, \hat{P}_{k}); \mathcal{G}_{\ell}^{+}) + \| h_{\ell} \|_{\infty;\Omega_{\ell}^{0}}^{2} \stackrel{!}{\leq} 2\varepsilon \end{aligned}$$

for any  $\varepsilon > 0$ . This can be seen as follows: By Lemma 3.5 we may first choose  $\ell$  large such that  $\|h_{\ell}\|_{\infty;\Omega_{\ell}^{0}}^{2} \leq \varepsilon$ . After fixing  $\ell$  the "point-wise" convergence of the indicators in Lemma 4.1 allows us then to choose a suitable  $k \geq \ell$  with  $\mathcal{E}_{k}^{2}((\hat{U}_{k}, \hat{Y}_{k}, \hat{P}_{k}); \mathcal{G}_{\ell}^{+}) \leq \varepsilon$ . This yields for any fixed  $v \in H^{2}(\Omega) \cap \mathring{H}^{1}(\Omega)$ 

$$\langle \mathcal{R}(\hat{y}_{\infty}; \hat{u}_{\infty}), v \rangle = \lim_{k \to \infty} \langle \mathcal{R}(\hat{Y}_k; \hat{U}_k), v \rangle = 0,$$

observing that  $\mathcal{R}$  is continuous with respect to its arguments and recalling the convergence  $(\hat{U}_k, \hat{Y}_k) \to (\hat{u}_\infty, \hat{y}_\infty)$  shown in Proposition 3.3. Since v is arbitrary we have shown  $\mathcal{R}(\hat{y}_\infty; \hat{u}_\infty) = 0$  in  $\mathbb{Y}^*$ . This in turn implies  $\hat{y}_\infty = S\hat{u}_\infty$  and finishes the proof.

**Convergence of error and estimator.** We are now in the position to prove the main result, where we again use the abbreviation  $\mathbb{W} = \mathbb{U} \times \mathbb{Y} \times \mathbb{Y}$ .

Proof of Theorem 1.2. Combining Propositions 2.1, 3.3, and 4.2 we obtain

$$\begin{split} \lim_{k \to \infty} \| (\hat{U}_k, \hat{Y}_k, \hat{P}_k) - (\hat{u}, \hat{p}, \hat{y}) \|_{\mathbb{W}} &\simeq \lim_{k \to \infty} \| (\hat{Y}_k, \hat{P}_k) - (S\hat{U}_k, S^* (\hat{Y}_k - y_d)) \|_{\mathbb{Y} \times \mathbb{Y}} \\ &= \| (\hat{y}_{\infty}, \hat{p}_{\infty}) - (S\hat{u}_{\infty}, S^* (\hat{y}_{\infty} - y_d) \|_{\mathbb{Y} \times \mathbb{Y}} = 0 \end{split}$$

This shows convergence of the error.

To show convergence of the estimator we decompose for  $k \geq \ell$  as in the proof to Proposition 4.2

$$\mathcal{E}_k^2((\hat{U}_k, \hat{Y}_k, \hat{P}_k); \mathcal{G}_k) = \mathcal{E}_k^2((\hat{U}_k, \hat{Y}_k, \hat{P}_k); \mathcal{G}_\ell^+) + \mathcal{E}_k^2((\hat{U}_k, \hat{Y}_k, \hat{P}_k); \mathcal{G}_k \setminus \mathcal{G}_\ell^+).$$

We first bound the second term on the right hand side. The local lower bound of Theorem 2.2 in combination with the finite overlap of the patches  $\mathcal{N}_k(E)$  allows us to bound

$$\begin{aligned} \mathcal{E}_{k}^{2}((\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k});\mathcal{G}_{k}\setminus\mathcal{G}_{\ell}^{+}) &\lesssim \|(\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k}) - (\hat{u},\hat{p},\hat{y})\|_{\mathbb{W}}^{2} + \sum_{E\in\mathcal{G}_{k}\setminus\mathcal{G}_{\ell}^{+}} \operatorname{osc}_{k}^{2}(\hat{U}_{k},y_{d};E) \\ &\lesssim \|(\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k}) - (\hat{u},\hat{p},\hat{y})\|_{\mathbb{W}}^{2} + \|h_{\ell}\|_{\infty;\Omega_{\ell}^{0}}^{2} \big(\|\hat{U}_{k}\|_{2;\Omega}^{2} + \|y_{d}\|_{2;\Omega}^{2}\big), \end{aligned}$$

using (3.3) and the rough estimate

$$\begin{aligned} \operatorname{psc}_{k}^{2}(\hat{U}_{k}, y_{d}; E) &= h_{E}^{2} \left( \| \hat{U}_{k} - \mathbb{P}_{\mathcal{G}_{k}} \hat{U}_{k} \|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^{2} + \| y_{d} - \mathbb{P}_{\mathcal{G}_{k}} y_{d} \|_{2;\Omega(\mathcal{N}_{\mathcal{G}}(E))}^{2} \right) \\ &\leq \| h_{\ell} \|_{\infty;\Omega_{\ell}^{0}}^{2} \left( \| \hat{U}_{k} \|_{2;\Omega(\mathcal{N}_{k}(E))}^{2} + \| y_{d} \|_{2;\Omega(\mathcal{N}_{k}(E))}^{2} \right). \end{aligned}$$

Since  $\|\hat{U}_k\|_{2;\Omega}^2 + \|y_d\|_{2;\Omega}^2 \lesssim 1$  we find

$$\mathcal{E}_{k}^{2}((\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k});\mathcal{G}_{k}) \lesssim \mathcal{E}_{k}^{2}((\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k});\mathcal{G}_{\ell}^{+}) + \|(\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k}) - (\hat{u},\hat{p},\hat{y})\|_{\mathbb{W}}^{2} + \|h_{\ell}\|_{\infty;\Omega_{\ell}^{0}}^{2}.$$

By Lemma 3.5 the last term  $\|h_{\ell}\|_{\infty;\Omega_{\ell}^{0}}^{2}$  can be made small by choosing  $\ell$  large. After fixing  $\ell$  we may choose as in the proof to Proposition 4.2 a  $k \geq \ell$  such that  $\mathcal{E}_{k}^{2}((\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k});\mathcal{G}_{\ell}^{+})$  is small. Moreover, the error convergence established above implies that the middle term  $\|(\hat{U}_{k},\hat{Y}_{k},\hat{P}_{k})-(\hat{u},\hat{p},\hat{y})\|_{\mathbb{W}}^{2}$  is small too, if we possibly increase k further. In summary, for any  $\varepsilon > 0$  we find a k such that

$$\mathcal{E}_k((\hat{U}_k, \hat{Y}_k, \hat{P}_k); \mathcal{G}_k) \leq \varepsilon.$$

This yields  $\mathcal{E}_k((\hat{U}_k, \hat{Y}_k, \hat{P}_k); \mathcal{G}_k) \to 0$  as  $k \to \infty$  and finishes the proof.

#### 5. Extensions and Outlook

The presented theory has been extended into several directions in the PhD thesis of the first author [6]. A detailed report is in preparation [7].

General linear-quadratic optimal control problem. The abstract framework can be found in [8, §2.1] and may be summarized as follows. We can allow for continuous, non-coercive bilinear forms  $\mathcal{B} \colon \mathbb{Y} \times \mathbb{Y} \to \mathbb{R}$  that satisfy an inf-sup condition.

This setting includes saddle point problems like the Stokes system and other mixed formulations. More general objectives  $\psi(y)$  can replace the simple tracking type functional  $\|y - y_d\|_{2;\Omega}^2$ . The functional  $\psi$  has to be quadratic and strictly convex. Its Fréchet-derivative  $\psi'$  has to satisfy a Lipschitz-condition. We may also consider any type of control space such that  $\mathbb{Y} \hookrightarrow \mathbb{U} \hookrightarrow \mathbb{Y}^*$  is a Gelfand triple. This then covers more general cases of distributed control as well as Neumann-boundary control. We are currently working on an extension for Dirichlet-boundary control [3].

Admitting a general class of PDE constraints requires appropriate assumptions on the estimators for the linear problems (2.1) and (2.2). Quite weak assumption are summarized in [16, §2.2.3] comprising other estimators like the hierarchical estimator, an estimator based on local problems on stars, an equilibrated residual estimator, and the ZZ-estimator; compare for instance with [10] for a detailed description of the diverse estimators. We may also weaken the assumption on marking to include marking strategies that adaptively focus on specific estimator contributions, like the indicators for the error in the state or adjoint equation. Such strategies are used in a comparison of adaptive strategies for optimal control problems in [8, §6]. We refer to [16, §2.2.4 and §5] for a sufficient and essentially necessary assumption on marking.

Most of the changes in the presented analysis are then concentrated in the proof to Lemma 3.2. This proof gets inevitably more involved due to the general structure of  $\psi$ , where one has to appropriately use convexity of  $\psi$ . All other statements can be proven using similar arguments with minor adjustments.

**Discretized control.** Up to now we have concentrated on the variational discretization of Hinze [5]. Here, the precise structure of the set of admissible controls  $\mathbb{U}^{\mathrm{ad}}$ is not of importance. The actual computation of a discrete solution yet requires the exact computation of  $\Pi(P)$  for a discrete function  $P \in \mathbb{Y}(\mathcal{G})$ . This typically gives restrictions on  $\mathbb{U}^{\mathrm{ad}}$ , like box-constraints with piecewise constant obstacles.

Very often the control space  $\mathbb{U}$  is discretized by a conforming finite element space  $\mathbb{U}(\mathcal{G})$ . Upon setting  $\mathbb{U}^{\mathrm{ad}}(\mathcal{G}) := \mathbb{U}^{\mathrm{ad}} \cap \mathbb{U}(\mathcal{G})$  and assuming that  $\mathbb{U}^{\mathrm{ad}}(\mathcal{G})$  is nonempty we can define a discrete projection operator  $\Pi_{\mathcal{G}} : \mathbb{U} \to \mathbb{U}^{\mathrm{ad}}(\mathcal{G})$  for  $p \in \mathbb{U}$ by

$$\Pi_{\mathcal{G}}(p) \in \mathbb{U}^{\mathrm{ad}}(\mathcal{G}): \qquad \langle \alpha \Pi_{\mathcal{G}}(p) + p, \, \Pi_{\mathcal{G}}(p) - U \rangle \leq 0 \qquad \forall U \in \mathbb{U}^{\mathrm{ad}}(\mathcal{G}).$$

An efficient computation of  $\Pi_{\mathcal{G}}$  benefits from a simple structure of  $\mathbb{U}^{\mathrm{ad}}$  and a suitable discrete control space  $\mathbb{U}(\mathcal{G})$ .

We can still consider the general setting of the previous paragraph. However, the analysis of adaptive finite elements for discretized controls gets painstakingly more laborious at several instances that we shortly list.

(1) The right hand side in the basic error equivalence in Proposition 2.1 has to be extended by the term  $\|\hat{U}_{\mathcal{G}} - \Pi(\hat{U}_{\mathcal{G}})\|_{\mathbb{U}}$  resulting in

$$\begin{aligned} \|(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\hat{u}, \hat{p}, \hat{y})\|_{\mathbb{W}} &\simeq \|(\hat{U}_{\mathcal{G}}, \hat{Y}_{\mathcal{G}}, \hat{P}_{\mathcal{G}}) - (\Pi(\hat{P}_{\mathcal{G}}), S(\hat{U}_{\mathcal{G}}), S^*\psi'(\hat{Y}_{\mathcal{G}}))\|_{\mathbb{W}};\\ \text{compare with [8, Theorem 2.2]} \end{aligned}$$

- (2) As a consequence, the element indicators of the estimator in Theorem 2.2 have to be enriched by a term  $\|\hat{U}_{\mathcal{G}} - \Pi(\hat{P}_{\mathcal{G}})\|^2_{\mathbb{U}(E)} = \|\Pi_{\mathcal{G}}(\hat{P}_{\mathcal{G}}) - \Pi(\hat{P}_{\mathcal{G}})\|^2_{\mathbb{U}(E)}$ to grant reliability of the estimator [8, Theorem 3.2]. Frequently this term is estimated further in order to completely avoid the computation of the continuous projection operator  $\Pi(\hat{P}_{\mathcal{G}})$  [4, 12]. This typically results in a nonefficient estimator; compare with [8, Remark 6.1].
- (3) Nesting of spaces  $\mathbb{Y}_k \subset \mathbb{Y}_{k+1}$  is essential to verify with current techniques the point-wise convergence of the solution operators  $S_k \to S_\infty$ ,  $S_k^* \to S_\infty^*$  in Proposition 3.1.

Likewise, for establishing the convergence  $\Pi_k(\hat{P}_k) \to \hat{u}_{\infty} = \Pi_{\infty}(\hat{p}_{\infty})$  in Lemma 3.2 nesting  $\mathbb{U}_k^{\mathrm{ad}} \subset \mathbb{U}_{k+1}^{\mathrm{ad}}$  of the sets of discrete admissible controls is instrumental. This nesting poses restrictions on data describing the set of admissible controls  $\mathbb{U}^{\mathrm{ad}}$ . Typically, such data has to be discrete over  $\mathcal{G}_0$ . In the proof to Lemma 3.2 we additionally have to account for the typical situation  $\hat{u}_{\infty} \notin \mathbb{U}_k^{\mathrm{ad}}$ . This increases substantially the complexity of the proof.

(4) The finite element spaces  $\{\mathbb{Y}_k\}_{k\geq 0}$  are "locally dense" in " $\Omega_{\infty}^0 := \lim_{k\to\infty} \Omega_k^0$ " in that  $\min_{V\in\mathbb{Y}_k} \|v-V\|_{\mathbb{Y}(\Omega_k^0)} \to 0$  as  $k\to\infty$ ; compare with [16, Remark 3.4]. Philosophically speaking, the improved bounds (2.8) and (2.9) for the residuals allow us to access this local density for showing that the limiting residuals  $\mathcal{R}(\hat{y}_{\infty}; \hat{u}_{\infty}), \mathcal{R}^*(\hat{p}_{\infty}; \psi'(\hat{y}_{\infty})) \in \mathbb{Y}^*$  are not supported in  $\Omega_{\infty}^0$ .

The additional contribution for the control error requires to establish the convergence

$$\lim_{k \to \infty} \|\hat{U}_k - \Pi(\hat{P}_k)\|_{\mathbb{U}(\Omega_k^0)} = \lim_{k \to \infty} \|\Pi_k(\hat{P}_k) - \Pi(\hat{P}_k)\|_{\mathbb{U}(\Omega_k^0)} \to 0.$$
(5.1)

For  $\mathbb{U} = L_2$  and piece-wise constant box-constraints in combination with a discontinuous or a continuous, piecewise linear control discretization one can verify (5.1) employing local density of  $\{\mathbb{U}_k\}_{k\geq 0}$  and point-wise properties of  $\Pi$ ; compare with [6, §8.4.2 and §8.4.3]. A characterization of properties of  $\Pi$  and  $\mathbb{U}(\mathcal{G})$  that ensure (5.1) is a challenging question and topic of future research.

(5) The proof of the estimator convergence in Theorem 1.2 strongly relies on local efficiency of the indicators as stated in Theorem 2.2. For discretized control this requires  $\|\hat{U}_{\mathcal{G}} - \Pi(\hat{P}_{\mathcal{G}})\|_{\mathbb{U}(E)}$  to be locally efficient, which can be shown if  $\Pi$  and  $\Pi_{\mathcal{G}}$  are locally Lipschitz continuous with uniformly bounded Lipschitz constants. This is typically true in case of distributed control.

In case of Neumann boundary control Lipschitz continuity of  $\Pi$  and  $\Pi_{\mathcal{G}}$ involves the trace operator  $T : H^1(\Omega) \to L_2(\partial\Omega)$ . We may therefore show global efficiency for  $\|\hat{U}_{\mathcal{G}} - \Pi(\hat{P}_{\mathcal{G}})\|_{L_2(\partial\Omega)}$  using the trace inequality on  $\Omega$ . An estimate of  $\|\hat{U}_{\mathcal{G}} - \Pi(\hat{P}_{\mathcal{G}})\|_{L_2(\partial E \cap \partial\Omega)}$  needs a local trace inequality on E. The typical scaling arguments yield negative powers of the local mesh-size and thereby ruling out local efficiency. As a consequence, we still can verify the error convergence of Theorem 1.2 but a proof of estimator convergence may require new techniques in that case.

#### References

- I. BABUŠKA AND M. VOGELIUS, Feedback and adaptive finite element solution of one-dimensional boundary value problems, Numer. Math., 44 (1984), pp. 75–102.
- [2] A. GAEVSKAYA, R. H. W. HOPPE, Y. ILIASH, AND M. KIEWEG, Convergence analysis of an adaptive finite element method for distributed control problems with control constraints, in Control of coupled partial differential equations, vol. 155 of Internat. Ser. Numer. Math., Birkhäuser, Basel, 2007, pp. 47–68.
- [3] F. D. GASPOZ AND K. G. SIEBERT, Adaptive finite elements for optimal control problems with Dirichlet boundary control. In Preparation.
- [4] M. HINTERMÜLLER, R. H. W. HOPPE, Y. ILIASH, AND M. KIEWEG, An a posteriori error analysis of adaptive finite element methods for distributed elliptic control problems with control constraints, ESAIM, Control Optim. Calc. Var., 14 (2008), pp. 540–560.
- [5] M. HINZE, A variational discretization concept in control constrained optimization: the linear-quadratic case, Comput. Optim. Appl., 30 (2005), pp. 45–61.
- [6] K. KOHLS, An Adaptive Finite Element Method for Control-Constrained Optimal Control Problems, PhD thesis, Fachbereich Mathematik, Unversität Stuttgart, 2013.
- [7] K. KOHLS, A. RÖSCH, AND K. G. SIEBERT, Convergence analysis of adaptive finite elements for optimal control problems. In Preparation.
- [8] —, A posteriori error analysis of optimal control problems with control constraints. Stuttgarter Mathematische Berichte 2013-001, Preprint SPP-1252-149.
- [9] K. KOHLS, A. RÖSCH, AND K. G. SIEBERT, A posteriori error estimators for control constrained optimal control problems, in Constrained Optimiziation and Optimal Control for Partial Differential Equations, L. et al., ed., International Series of Numerical Mathematics 160, Springer, 2012, pp. 431–443.
- [10] C. KREUZER AND K. G. SIEBERT, Decay rates of adaptive finite elements with Dörfler marking, Numer. Math., 117 (2011), pp. 679–716.
- [11] J. L. LIONS, Optimal Control of Systems Governed by Partial Differential Equations, vol. 170 of Die Grundlehren der mathematischen Wissenschaften, Springer-Verlag, 1971.
- [12] W. LIU AND N. YAN, A posteriori error estimates for distributed convex optimal control problems, Adv. Comput. Math, 15 (2001), pp. 285–309.
- [13] C. MAKRIDAKIS AND R. H. NOCHETTO, Elliptic reconstruction and a posteriori error estimates for parabolic problems, SIAM J. Numer. Anal., 41 (2003), pp. 1585–1594.
- [14] P. MORIN, K. G. SIEBERT, AND A. VEESER, A basic convergence result for conforming adaptive finite elements, Math. Models Methods Appl. Sci., 18 (2008), pp. 707– 737.
- [15] A. SCHMIDT AND K. G. SIEBERT, Design of adaptive finite element software, vol. 42 of Lecture Notes in Computational Science and Engineering, Springer-Verlag, Berlin, 2005. The finite element toolbox ALBERTA, With 1 CD-ROM (Unix/Linux).
- [16] K. G. SIEBERT, A convergence proof for adaptive finite elements without lower bound., IMA J. Numer. Anal., 31 (2011), pp. 947–970.
- [17] F. TRÖLTZSCH, Optimal control of partial differential equations, vol. 112 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2010.

Theory, methods and applications, Translated from the 2005 German original by Jürgen Sprekels.

Kristina Kohls Institut für Angewandte Analysis und Numerische Simulation, Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, D-70569 Stuttgart, Germany URL: http://www.ians.uni-stuttgart.de/nmh/ e-mail: kristina.kohls@ians.uni-stuttgart.de

Arnd Rösch Fakultät für Mathematik, Universität Duisburg-Essen, Forsthausweg 2, D-47057 Duisburg, Germany URL: http://www.uni-due.de/mathematik/agroesch/ e-mail: arnd.roesch@uni-due.de

Kunibert G. Siebert Institut für Angewandte Analysis und Numerische Simulation, Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, D-70569 Stuttgart, Germany URL: http://www.ians.uni-stuttgart.de/nmh/ e-mail: kg.siebert@ians.uni-stuttgart.de

#### Erschienene Preprints ab Nummer 2007/2007-001

Komplette Liste: http://www.mathematik.uni-stuttgart.de/preprints

- 2013-011 Kohls, K; Rösch, A.; Siebert, K.G.: Convergence of Adaptive Finite Elements for Control Constrained Optimal Control Problems
- 2013-010 *Corli, A.; Rohde, C.; Schleper, V.:* Parabolic Approximations of Diffusive-Dispersive Equations
- 2013-009 Nava-Yazdani, E.; Polthier, K.: De Casteljau's Algorithm on Manifolds
- 2013-008 *Bächle, A.; Margolis, L.:* Rational conjugacy of torsion units in integral group rings of non-solvable groups
- 2013-007 Knarr, N.; Stroppel, M.J.: Heisenberg groups over composition algebras
- 2013-006 Knarr, N.; Stroppel, M.J.: Heisenberg groups, semifields, and translation planes
- 2013-005 *Eck, C.; Kutter, M.; Sändig, A.-M.; Rohde, C.:* A Two Scale Model for Liquid Phase Epitaxy with Elasticity: An Iterative Procedure
- 2013-004 Griesemer, M.; Wellig, D.: The Strong-Coupling Polaron in Electromagnetic Fields
- 2013-003 *Kabil, B.; Rohde, C.:* The Influence of Surface Tension and Configurational Forces on the Stability of Liquid-Vapor Interfaces
- 2013-002 *Devroye, L.; Ferrario, P.G.; Györfi, L.; Walk, H.:* Strong universal consistent estimate of the minimum mean squared error
- 2013-001 *Kohls, K.; Rösch, A.; Siebert, K.G.:* A Posteriori Error Analysis of Optimal Control Problems with Control Constraints
- 2012-018 *Kimmerle, W.; Konovalov, A.:* On the Prime Graph of the Unit Group of Integral Group Rings of Finite Groups II
- 2012-017 *Stroppel, B.; Stroppel, M.:* Desargues, Doily, Dualities, and Exceptional Isomorphisms
- 2012-016 *Moroianu, A.; Pilca, M.; Semmelmann, U.:* Homogeneous almost guaternion-Hermitian manifolds
- 2012-015 *Steinke, G.F.; Stroppel, M.J.:* Simple groups acting two-transitively on the set of generators of a finite elation Laguerre plane
- 2012-014 *Steinke, G.F.; Stroppel, M.J.:* Finite elation Laguerre planes admitting a two-transitive group on their set of generators
- 2012-013 *Diaz Ramos, J.C.; Dominguez Vázquez, M.; Kollross, A.:* Polar actions on complex hyperbolic spaces
- 2012-012 Moroianu; A.; Semmelmann, U.: Weakly complex homogeneous spaces
- 2012-011 Moroianu; A.; Semmelmann, U.: Invariant four-forms and symmetric pairs
- 2012-010 Hamilton, M.J.D.: The closure of the symplectic cone of elliptic surfaces
- 2012-009 Hamilton, M.J.D.: Iterated fibre sums of algebraic Lefschetz fibrations
- 2012-008 Hamilton, M.J.D.: The minimal genus problem for elliptic surfaces
- 2012-007 *Ferrario, P.:* Partitioning estimation of local variance based on nearest neighbors under censoring
- 2012-006 Stroppel, M.: Buttons, Holes and Loops of String: Lacing the Doily
- 2012-005 Hantsch, F.: Existence of Minimizers in Restricted Hartree-Fock Theory
- 2012-004 Grundhöfer, T.; Stroppel, M.; Van Maldeghem, H.: Unitals admitting all translations
- 2012-003 Hamilton, M.J.D.: Representing homology classes by symplectic surfaces
- 2012-002 Hamilton, M.J.D.: On certain exotic 4-manifolds of Akhmedov and Park

- 2012-001 Jentsch, T.: Parallel submanifolds of the real 2-Grassmannian
- 2011-028 Spreer, J.: Combinatorial 3-manifolds with cyclic automorphism group
- 2011-027 *Griesemer, M.; Hantsch, F.; Wellig, D.:* On the Magnetic Pekar Functional and the Existence of Bipolarons
- 2011-026 Müller, S.: Bootstrapping for Bandwidth Selection in Functional Data Regression
- 2011-025 *Felber, T.; Jones, D.; Kohler, M.; Walk, H.:* Weakly universally consistent static forecasting of stationary and ergodic time series via local averaging and least squares estimates
- 2011-024 Jones, D.; Kohler, M.; Walk, H.: Weakly universally consistent forecasting of stationary and ergodic time series
- 2011-023 *Györfi, L.; Walk, H.:* Strongly consistent nonparametric tests of conditional independence
- 2011-022 *Ferrario, P.G.; Walk, H.:* Nonparametric partitioning estimation of residual and local variance based on first and second nearest neighbors
- 2011-021 Eberts, M.; Steinwart, I.: Optimal regression rates for SVMs using Gaussian kernels
- 2011-020 Frank, R.L.; Geisinger, L.: Refined Semiclassical Asymptotics for Fractional Powers of the Laplace Operator
- 2011-019 *Frank, R.L.; Geisinger, L.:* Two-term spectral asymptotics for the Dirichlet Laplacian on a bounded domain
- 2011-018 Hänel, A.; Schulz, C.; Wirth, J.: Embedded eigenvalues for the elastic strip with cracks
- 2011-017 Wirth, J.: Thermo-elasticity for anisotropic media in higher dimensions
- 2011-016 Höllig, K.; Hörner, J.: Programming Multigrid Methods with B-Splines
- 2011-015 *Ferrario, P.:* Nonparametric Local Averaging Estimation of the Local Variance Function
- 2011-014 *Müller, S.; Dippon, J.:* k-NN Kernel Estimate for Nonparametric Functional Regression in Time Series Analysis
- 2011-013 Knarr, N.; Stroppel, M.: Unitals over composition algebras
- 2011-012 *Knarr, N.; Stroppel, M.:* Baer involutions and polarities in Moufang planes of characteristic two
- 2011-011 Knarr, N.; Stroppel, M.: Polarities and planar collineations of Moufang planes
- 2011-010 Jentsch, T.; Moroianu, A.; Semmelmann, U.: Extrinsic hyperspheres in manifolds with special holonomy
- 2011-009 *Wirth, J.:* Asymptotic Behaviour of Solutions to Hyperbolic Partial Differential Equations
- 2011-008 Stroppel, M.: Orthogonal polar spaces and unitals
- 2011-007 *Nagl, M.:* Charakterisierung der Symmetrischen Gruppen durch ihre komplexe Gruppenalgebra
- 2011-006 *Solanes, G.; Teufel, E.:* Horo-tightness and total (absolute) curvatures in hyperbolic spaces
- 2011-005 Ginoux, N.; Semmelmann, U.: Imaginary Kählerian Killing spinors I
- 2011-004 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic *D*-Scales: Part II Gain-Scheduled Control
- 2011-003 *Scherer, C.W.; Köse, I.E.:* Control Synthesis using Dynamic *D*-Scales: Part I Robust Control

- 2011-002 Alexandrov, B.; Semmelmann, U.: Deformations of nearly parallel G2-structures
- 2011-001 Geisinger, L.; Weidl, T.: Sharp spectral estimates in domains of infinite volume
- 2010-018 Kimmerle, W.; Konovalov, A.: On integral-like units of modular group rings
- 2010-017 *Gauduchon, P.; Moroianu, A.; Semmelmann, U.:* Almost complex structures on quaternion-Kähler manifolds and inner symmetric spaces
- 2010-016 Moroianu, A.; Semmelmann, U.: Clifford structures on Riemannian manifolds
- 2010-015 *Grafarend, E.W.; Kühnel, W.:* A minimal atlas for the rotation group SO(3)
- 2010-014 Weidl, T.: Semiclassical Spectral Bounds and Beyond
- 2010-013 Stroppel, M.: Early explicit examples of non-desarguesian plane geometries
- 2010-012 Effenberger, F.: Stacked polytopes and tight triangulations of manifolds
- 2010-011 *Györfi, L.; Walk, H.:* Empirical portfolio selection strategies with proportional transaction costs
- 2010-010 *Kohler, M.; Krzyżak, A.; Walk, H.:* Estimation of the essential supremum of a regression function
- 2010-009 *Geisinger, L.; Laptev, A.; Weidl, T.:* Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010-008 Poppitz, S.; Stroppel, M.: Polarities of Schellhammer Planes
- 2010-007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.:* Non-existence of isomorphisms between certain unitals
- 2010-006 *Höllig, K.; Hörner, J.; Hoffacker, A.:* Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods
- 2010-005 Kaltenbacher, B.; Walk, H.: On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010-004 Kühnel, W.; Solanes, G.: Tight surfaces with boundary
- 2010-003 *Kohler, M; Walk, H.:* On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010-002 *Gulde, M.; Stroppel, M.:* Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010-001 *Leitner, F.:* Examples of almost Einstein structures on products and in cohomogeneity one
- 2009-008 Griesemer, M.; Zenk, H.: On the atomic photoeffect in non-relativistic QED
- 2009-007 *Griesemer, M.; Moeller, J.S.:* Bounds on the minimal energy of translation invariant n-polaron systems
- 2009-006 *Demirel, S.; Harrell II, E.M.:* On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009-005 Bächle, A, Kimmerle, W.: Torsion subgroups in integral group rings of finite groups
- 2009-004 Geisinger, L.; Weidl, T.: Universal bounds for traces of the Dirichlet Laplace operator
- 2009-003 Walk, H.: Strong laws of large numbers and nonparametric estimation
- 2009-002 Leitner, F.: The collapsing sphere product of Poincaré-Einstein spaces
- 2009-001 Brehm, U.; Kühnel, W.: Lattice triangulations of  $E^3$  and of the 3-torus
- 2008-006 *Kohler, M.; Krzyżak, A.; Walk, H.:* Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps

- 2008-005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.:* Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems
- 2008-004 *Leitner, F.:* Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008-003 Effenberger, F.; Kühnel, W.: Hamiltonian submanifolds of regular polytope
- 2008-002 *Hertweck, M.; Höfert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups PSL(2, q)
- 2008-001 *Kovarik, H.; Vugalter, S.; Weidl, T.:* Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007-006 Weidl, T .: Improved Berezin-Li-Yau inequalities with a remainder term
- 2007-005 *Frank, R.L.; Loss, M.; Weidl, T.:* Polya's conjecture in the presence of a constant magnetic field
- 2007-004 Ekholm, T.; Frank, R.L.; Kovarik, H.: Eigenvalue estimates for Schrödinger operators on metric trees
- 2007-003 Lesky, P.H.; Racke, R.: Elastic and electro-magnetic waves in infinite waveguides
- 2007-002 Teufel, E.: Spherical transforms and Radon transforms in Moebius geometry
- 2007-001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions