Universität Stuttgart

Fachbereich Mathematik

On the measure of Voronoi cells

Luc Devroye, László Györfi, Gábor Lugosi, Harro Walk

Preprint 2016/006

Fachbereich Mathematik Fakultät Mathematik und Physik Universität Stuttgart Pfaffenwaldring 57 D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de
WWW: http://www.mathematik.uni-stuttgart.de/preprints

ISSN 1613-8309

C Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors. LaTEX-Style: Winfried Geis, Thomas Merkle

On the measure of Voronoi cells *

Luc Devroye[†] László Györfi[‡] Gábor Lugosi[§] Harro Walk[¶]

June 30, 2016

Abstract

We study the measure of a typical cell in a Voronoi tessellation defined by nindependent random points drawn from a density f in \mathbb{R}^d . In particular, we prove that the asymptotic distribution of the measure—with respect to $d\mu = f(x)dx$ —of the cell centered at a point $x \in \mathbb{R}^d$ is independent of x and the density f. We determine all moments of the asymptotic distribution and show that the distribution becomes more concentrated as d becomes large. In particular, we show that the variance converges to zero exponentially fast in d. We also obtain a bound independent of the density for the rate of convergence of the diameter of a typical Voronoi cell.

 ${\bf Keywords:} \ {\rm random} \ {\rm pointset}, \ {\rm Voronoi} \ {\rm tessellation}, \ {\rm stochastic} \ {\rm geometry}$

2010 Mathematics Subject Classification: 60D05

^{*}Luc Devroye was supported by the Natural Sciences and Engineering Research Council (NSERC) of Canada. László Györfi was supported in part by the National Development Agency (NFÜ, Hungary) as part of the project Introduction of Cognitive Methods for UAV Collision Avoidance Using Millimeter Wave Radar, grant KMR-12-1-2012-0008. Gábor Lugosi was supported by the Spanish Ministry of Economy and Competitiveness, Grant MTM2015-67304-P and FEDER, UE.

[†]McGill University, lucdevroye@gmail.com

[‡]Budapest University of Technology and Economics, gyorfi@cs.bme.hu

ICREA and Pompeu Fabra University, gabor.lugosi@upf.edu

[¶]Universität Stuttgart, harro.walk@t-online.de

1 Introduction

Let X_1, \ldots, X_n be independent, identically distributed random vectors taking values in \mathbb{R}^d . We denote the common distribution of the X_i by μ . We assume throughout the paper that μ is absolutely continuous with respect to the Lebesgue measure λ and denote the density of μ by f. Hence, $\mu(A) = \int_A f(x) dx$ for all Lebesgue measurable sets $A \subset \mathbb{R}^d$.

The X_i define the random subsets S_1, \ldots, S_n of \mathbb{R}^d such that S_i contains all points in \mathbb{R}^d that are strictly closer to X_i than any other point in $\{X_1, \ldots, X_n\} \setminus \{X_i\}$. Formally,

$$S_i = \left\{ x \in \mathbb{R}^d : \|x - X_i\| < \min_{j = \{1, \dots, n\} \setminus \{i\}} \|x - X_j\| \right\} .$$

 $\{S_1, \ldots, S_n\}$ is a so-called *Voronoi tessellation* and the S_i are the *Voronoi cells*. Here $\|\cdot\|$ denotes the usual Euclidean norm.

In this paper we are interested in the measure of a "typical" Voronoi cell. In particular, we study the conditional distribution of the random variable $\mu(S_1)$ conditioned on the event that $X_1 = x$ for some x in the support of μ . Equivalently, we study $M_n(x)$ where for any $x \in \mathbb{R}^d$, $M_n(x)$ is the μ -measure of the Voronoi cell S_1 with nucleus x associated with the point process $\{x, X_2, \ldots, X_n\}$.

Note that since $\sum_{j=1}^{n} \mu(S_j) = 1$ and $\mu(S_1), \ldots, \mu(S_n)$ are identically distributed, we have $n\mathbb{E}\left[\mu(S_1)\right] = 1$. In Theorem 1 below we prove that, for μ -almost all x, we have $n\mathbb{E}M_n(x) \to 1$. We also show that $n^2\mathbb{E}[M_n(x)^2]$ converges to a limit that is independent of x and the distribution μ . In fact, we prove that for μ -almost all x, $nM_n(x)$ has a limiting distribution that only depends on the dimension. We show that the limiting distribution becomes more concentrated as the dimension d grows. In particular, the variance converges to zero exponentially fast in d.

Finally, we study the diameter diam (S_1) of the Voronoi cell centered at X_1 . We show that for μ -almost all x, conditional on $X_1 = x$, diam (S_1) converges to zero at a rate of $n^{-1/d}$.

Throughout the paper, $B_{x,r}$ denotes the closed ball of radius r > 0 centered at $x \in \mathbb{R}^d$.

Motivation and related work

Nearest-neighbor-based estimators are among the most basic and popular tools in addressing classification, regression, density estimation and other problems in nonparametric statistics, see Biau and Devroye [2] for a recent survey. Many of these estimators build on a Voronoi partition of the space based on data drawn from an unknown density. In statistics it is important to understand the behavior of the estimators under as few assumptions as possible on the distribution of the data. In particular, the measure and the diameter of the Voronoi cells give valuable information about the properties of nearest-neighbor estimators. The distribution-free results derived in this paper were motivated by and contribute to the understanding of such estimators.

The measure of a "typical" cell in a Voronoi tessellation has been mostly studied in the case when the points are drawn from a homogeneous Poisson process. Asymptotically, as $n \to \infty$, this is equivalent to the special case of drawing n points from the uniform distribution μ on (say) the unit ball. The study of the (Lebesgue) measure of Voronoi cells dates back to at least Gilbert [7] who derived formulas and numerical estimates for the second and third moments the measure of a Voronoi cell when d = 2 or 3. See also Brakke [3, 4], Hayen and Quine [8].

Our notion of the distribution of a typical cell is analogous to the so-called "Palm distribution" of the volume of a Voronoi cell in stochastic geometry—Chiu, Stoyan, Kendall, and Mecke [5], Møller [12], Møller and Stoyan [13].

Brakke [3, 4], Hayen and Quine [8], Heinrich et al. [9], Heinrich and Muche [10], Zuyev [16] study characteristics of "typical" cells in a Voronoi tessallation of a homogeneous Poisson process, including the second moment of the volume.

The results of the paper may be easily extended to inhomogeneous multi-dimensional Poisson processes. This extension is relevant for modelling mobile communication networks, see Baccelli et al. [1].

For a survey and comprehensive treatment of Voronoi tessellations, we refer to Okabe, Boots, Sugihara, and Chiu [14].

2 Convergence of the first two moments

Theorem 1 below establishes the asymptotic value of the first and second moments of the measure of a typical cell centered at a point x. The remarkable feature is that the asymptotic values are independent of both the density f and the point x (for μ -almost all x) and only depend on the dimension d. In fact, in Theorem 2 we show that the limit distribution is also independent of f and x. We emphasize that both theorems hold without any assumption on the density f.

The asymptotic second moment is expressed in terms of a random variable W defined as follows. Let Y be a random vector uniformly distributed in $B_{0,1}$. Define $\overline{1} = (1, 0, 0, \dots, 0) \in \mathbb{R}^d$ and let $\overline{B} = B_{\overline{1},1} \bigcup B_{Y,||Y||}$. Introduce the random variable

$$W = \frac{\lambda(\overline{B})}{\lambda(B_{0,1})} \tag{1}$$

(recall that λ denotes the Lebesgue measure) and let

$$\alpha(d) \stackrel{\text{def}}{=} \mathbb{E}\left[\frac{2}{W^2}\right]$$

Theorem 1 Let μ have a density f. Then, as $n \to \infty$,

(i)

$$n\mathbb{E}M_n(x) \to 1$$
 for μ -almost all x .

(ii)

$$n^2 \mathbb{E}[M_n(x)^2] \to \alpha(d) \quad \text{for } \mu\text{-almost all } x$$
.

In Section 4 we obtain estimates for the asymptotic conditional second moment $\alpha(d)$. In particular, in Theorem 3 we show that for all dimensions, $1 \leq \alpha(d) \leq 1 + 6(3/4)^{d/2}$ and therefore the asymptotic variance of $\mu(S_1)$ (conditioned on $X_1 = x$) decreases to zero exponentially in d.

Lebesgue density theorem. The proofs use a version of the Lebesgue density theorem that we recall first.

Let \mathcal{B} be the class of all closed balls of \mathbb{R}^d containing the origin. We say that x is a *Lebesgue point* for f if for all sequences $B_k \in \mathcal{B}$ with $\lambda(B_k) \downarrow 0$,

$$\lim_{k \to \infty} \frac{\int_{x+B_k} f(y) dy}{\lambda(B_k)} = f(x) \; .$$

Let A be the set of all $x \in \mathbb{R}^d$ such that f(x) > 0 and x is a Lebesgue point for f. Then $\mu(A) = 1$ by Wheeden and Zygmund [15, pp. 106–108]. See also Devroye and Györfi [6], Chapter 2.

Proof of part (i). Observe that

$$\mathbb{E}M_n(x) = \mathbb{P}\left\{X_{n+1} \in S_1 \mid X_1 = x\right\} \\ = \mathbb{P}\left\{\bigcap_{i=2}^n \{X_i \notin B_{X_{n+1}, \|X_{n+1} - x\|}\}\right\} \\ = \mathbb{E}\left[(1 - Z(x))^{n-1}\right] ,$$

where $Z(x) = \mu(B_{X,||X-x||})$. (Recall that both X and X_{n+1} are independent of X_1, \ldots, X_n and have the same distribution.) First note that

$$n\mathbb{E}\left\{(1-Z(x))^{n-1}\right\} \to 1$$

whenever

$$\lim_{z \downarrow 0} z^{-1} \mathbb{P}\left\{ \mu(B_{X, \|X-x\|}) \le z \right\} = 1 .$$
 (2)

To see this, suppose that (2) holds and for every $\epsilon > 0$ there is a $\delta > 0$ such that $|\mathbb{P}\{Z(x) \leq z\} - z| < z\epsilon$ whenever $z \in (0, \delta)$. But then

$$\begin{split} n \mathbb{E} \left\{ (1 - Z(x))^{n-1} \right\} \\ &= n \int_0^1 \mathbb{P} \left\{ (1 - Z(x))^{n-1} \ge t \right\} dt \\ &= n(n-1) \int_0^1 \mathbb{P} \left\{ Z(x) \le z \right\} (1-z)^{n-2} dz \\ &\quad \text{(by a change of variables)} \\ &= n(n-1) \int_0^\delta \mathbb{P} \left\{ Z(x) \le z \right\} (1-z)^{n-2} dz + n(n-1) \int_0^\delta \mathbb{P} \left\{ Z(x) \le z \right\} (1-z)^{n-2} dz \\ &= n(n-1) \int_0^\delta z(1+\rho(z)) \mathbb{P} \left\{ Z(x) \le z \right\} (1-z)^{n-2} dz + O(n^2(1-\delta)^n) \\ &\quad \text{(where } |\rho(z)| < \epsilon) \\ &= 1 + O(\epsilon) + O(n^2(1-\delta)^n) \;. \end{split}$$

The intuitive reason of why (2) should hold is that for any x, $\mu(B_{x,\|X-x\|})$ is uniformly distributed on [0, 1] and that $\mu(B_{X,\|X-x\|})$ is stochastically close to $\mu(B_{x,\|X-x\|})$ when $\|X-x\|$ is small in a sense specified below. The rest of the proof establishes the convergence (2).

By the Lebesgue density theorem, it suffices to prove (2) for all Lebesgue points x with f(x) > 0. Fix such a point x. Since for any sequence $B_k \in \mathcal{B}$ with $\lambda(B_k) \downarrow 0$ we have

$$\frac{\mu(x+B_k)}{\lambda(x+B_k)} \to f(x) \; ,$$

for any $\epsilon \in (0, 1)$ we can find $\delta > 0$ (possibly depending on x) such that $||v - x|| \le \delta$ implies

$$\left|\frac{\mu(B_{v,\|v-x\|})}{\lambda(B_{v,\|v-x\|})} - f(x)\right| \le \epsilon f(x)$$

and

$$\left|\frac{\mu(B_{x,\|v-x\|})}{\lambda(B_{x,\|v-x\|})} - f(x)\right| \le \epsilon f(x) \ .$$

This also implies that for any v with $||v - x|| \ge \delta$,

$$\mu(B_{v,\|v-x\|}) \geq \mu(B_{v^*,\delta}) \geq (1-\epsilon)f(x)\lambda(B_{0,\delta})$$
and
$$\mu(B_{x,\|v-x\|}) \geq \mu(B_{x,\delta}) \geq (1-\epsilon)f(x)\lambda(B_{0,\delta})$$
(3)

where v^* is the unique point on the surface of $B_{x,\delta}$ and on the line segment (x, v]. Take z > 0 so small that $z < (1 - \epsilon)f(x)\lambda(B_{0,\delta})$. We again use the fact that $\mu(B_{x,||X-x||}) = U$ has the uniform distribution on [0, 1]. We rewrite

$$\mu(B_{X,\|X-x\|}) = \frac{\mu(B_{X,\|X-x\|})}{\lambda(B_{X,\|X-x\|})} \cdot \frac{\lambda(B_{x,\|X-x\|})}{\mu(B_{x,\|X-x\|})} \cdot \mu(B_{x,\|X-x\|}) .$$

If $||X - x|| \leq \delta$, then the product of the first two factors is sandwiched between

$$\frac{1-\epsilon}{1+\epsilon}$$
 and $\frac{1+\epsilon}{1-\epsilon}$.

By (3), we have $||X - x|| \leq \delta$ if either $\mu(B_{X,||X-x||}) \leq z$ or $\mu(B_{x,||X-x||}) \leq z$, and in that case,

$$\mu(B_{X,\|X-x\|}) \ge \frac{1-\epsilon}{1+\epsilon} \mu(B_{x,\|X-x\|})$$

and

$$\mu(B_{X,\|X-x\|}) \le \frac{1+\epsilon}{1-\epsilon}\mu(B_{x,\|X-x\|})$$

Thus,

$$\mathbb{P}\{\mu(B_{X,\|X-x\|}) \le z\} \le \mathbb{P}\left\{\frac{1-\epsilon}{1+\epsilon}\mu(B_{x,\|X-x\|}) \le z\right\}$$
$$\le \mathbb{P}\left\{\frac{1-\epsilon}{1+\epsilon}U \le z\right\}$$
$$= \min\left\{z\frac{1+\epsilon}{1-\epsilon}, 1\right\}.$$

Similarly, for $0 < z \le 1$,

$$\mathbb{P}\{\mu(B_{X,\|X-x\|}) \leq z\} = \mathbb{P}\{\mu(B_{X,\|X-x\|}) \leq z, \|X-x\| \leq \delta\}$$

$$\geq \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon}\mu(B_{x,\|X-x\|}) \leq z, \|X-x\| \leq \delta\right\}$$

$$= \mathbb{P}\left\{\mu(B_{x,\|X-x\|}) \leq \frac{1-\epsilon}{1+\epsilon}z, \|X-x\| \leq \delta\right\}$$

$$= \mathbb{P}\left\{\mu(B_{x,\|X-x\|}) \leq \frac{1-\epsilon}{1+\epsilon}z\right\}$$

$$= \mathbb{P}\left\{U \leq \frac{1-\epsilon}{1+\epsilon}z\right\}$$

$$= \frac{1-\epsilon}{1+\epsilon}z.$$

This proves (2) and part (i) of Theorem 1.

Proof of part (ii). Similarly to the proof of part (i), observe that

$$\mathbb{E}[M_n(x)^2] = \mathbb{P}\left\{X_{n+1} \in S_1, X_{n+2} \in S_1 \mid X_1 = x\right\}$$

= $\mathbb{P}\left\{\bigcap_{i=2}^n \{X_i \notin B_{X_{n+1}, \|X_{n+1} - x\|} \cup B_{X_{n+2}, \|X_{n+2} - x\|}\}\right\}$
= $\mathbb{E}\left[(1 - Z_2(x))^{n-1}\right]$,

where

$$Z_2(x) = \mu(B_{X, \|X-x\|} \cup B_{X', \|X'-x\|})$$

with X and X' independent and distributed with respect to μ . In analogy with the argument of part (i), in order to prove that

$$\lim_{n \to \infty} n^2 \mathbb{E} \left\{ (1 - Z_2(x))^{n-1} \right\} = \alpha(d) ,$$

it suffices to show that, for μ -almost all x,

$$\lim_{z \downarrow 0} z^{-2} \mathbb{P}\left\{ \mu(B_{X, \|X-x\|} \cup B_{X', \|X'-x\|}) \le z \right\} = \alpha(d)/2.$$

The rough idea of the proof is as follows. The approximate equalities are made rigorous

below. For small z,

$$\begin{aligned} z^{-2} \mathbb{P} \left\{ \mu(B_{X,\|X-x\|} \cup B_{X',\|X'-x\|}) \leq z \right\} \\ &= z^{-2} \mathbb{P} \left\{ \frac{\mu(B_{X,\|X-x\|} \cup B_{X',\|X'-x\|})}{\max\{\mu(B_{X,\|X-x\|}), \mu(B_{X',\|X'-x\|})\}} \max\{\mu(B_{X,\|X-x\|}), \mu(B_{X',\|X'-x\|})\} \leq z \right\} \\ &\underset{z \to 0}{\sim} z^{-2} \mathbb{P} \left\{ \frac{\mu(B_{X,\|X-x\|} \cup B_{X',\|X'-x\|})}{\max\{\mu(B_{X,\|X-x\|}), \mu(B_{X',\|X'-x\|})\}} \max\{\mu(B_{x,\|X-x\|}), \mu(B_{x,\|X'-x\|})\} \leq z \right\} \\ &\underset{z \to 0}{\sim} z^{-2} \mathbb{P} \left\{ W \max\{U_1, U_2\} \leq z \right\} \\ &\qquad (\text{where } U_1, U_2 \text{ are i.i.d. uniform, independent of } W) \\ &= z^{-2} \mathbb{P} \left\{ W U^{1/2} \leq z \right\} \\ &= z^{-2} \mathbb{E} \left[\min(z^2/W^2, 1) \right] \underset{z \to 0}{\sim} \mathbb{E} \left[\frac{1}{W^2} \right] . \end{aligned}$$

To prove the desired limit formally, as before, by the Lebesgue density theorem, we may assume that $x \in \mathbb{R}^d$ is such that f(x) > 0 and x is a Lebesgue point for f. A key point of the proof uses coupling. Let (Y_1, Y_2) be the canonical reordering of (X, X') such that

$$||Y_2 - x|| \ge ||Y_1 - x||$$
.

and introduce $M = \max(||X - x||, ||X' - x||) = ||Y_2 - x||$. Define the random variable N by

$$N = \begin{cases} 1 & \text{if } Y_1 = X' \\ 2 & \text{if } Y_2 = X' \end{cases}$$

Then set $V_2 = Y_2$ and let V_1 be uniformly distributed on $B_{x,\|V_2-x\|}$ such that V_1 is maximally coupled with Y_1 given Y_2 . From Doeblin's coupling argument,

$$\mathbb{P}\{Y_1 \neq V_1 \mid Y_2\} = \frac{1}{2} \int |f_{Y_1}(v) - f_{V_1}(v)| dv ,$$

where f_{Y_1} , f_{V_1} are the conditional densities of Y_1 and V_1 given Y_2 .

Choose $\delta > 0$ so small that for $M \leq \delta$, we have, simultaneously,

$$\frac{\mu(B_{x,M})}{\lambda(B_{x,M})} \in [f(x)(1-\epsilon), f(x)(1+\epsilon)] ,$$
$$\frac{\mu(B_{X,\|X-x\|} \cup B_{X',\|X'-x\|})}{\lambda(B_{X,\|X-x\|} \cup B_{X',\|X'-x\|})} \in [f(x)(1-\epsilon), f(x)(1+\epsilon)] ,$$

and

$$\frac{\mu(B_{X,M})}{\lambda(B_{X,M})} \left| \frac{\lambda(B_{x,M})}{\mu(B_{x,M})} - \frac{1}{f(x)} \right| + \frac{1}{\lambda(B_{x,M})f(x)} \int_{B_{x,M}} |f(v) - f(x)| dv \le \epsilon .$$

Such a δ exists by three applications of the Lebesgue density theorem. (Recall that x is a Lebesgue point.) Since

$$f_{Y_1}(v) = \frac{f(v)}{\mu(B_{x,\|Y_2 - x\|})} \mathbb{1}_{v \in B_{x,M}} \quad \text{and} \quad f_{V_1}(v) = \frac{1}{\lambda(B_{x,M})} \mathbb{1}_{v \in B_{x,M}} ,$$

(where $\mathbb{1}$ denotes the indicator function) we have, writing $B = B_{x,M}$,

$$\int |f_{Y_1}(v) - f_{V_1}(v)| dv = \int_B \left| \frac{f(v)}{\lambda(B)} \frac{\lambda(B)}{\mu(B)} - \frac{1}{\lambda(B)} \right| dv$$

$$\leq \frac{1}{\lambda(B)} \int_B f(v) \left| \frac{\lambda(B)}{\mu(B)} - \frac{1}{f(x)} \right| dv + \frac{1}{\lambda(B)} \int_B \left| \frac{f(v)}{f(x)} - 1 \right| dv$$

$$= \frac{\mu(B)}{\lambda(B)} \left| \frac{\lambda(B)}{\mu(B)} - \frac{1}{f(x)} \right| + \frac{1}{f(x)} \frac{1}{\lambda(B)} \int_B |f(v) - f(x)| dv$$

$$\leq \epsilon$$

if $M \leq \delta$, by choice of δ . Finally, define a pair of random variables (V, V'), both taking values in \mathbb{R}^d , as follows.

$$(V, V') = \begin{cases} (V_1, V_2) & \text{if } N = 2\\ (V_2, V_1) & \text{if } N = 1 \end{cases}$$

The argument above implies that

$$\mathbb{P}\{(V,V') \neq (X,X') \mid M\} \le \mathbb{1}_{M > \delta} + \mathbb{1}_{M \le \delta} \frac{\epsilon}{2} .$$

Since

$$(\mu(B_{x,\|X-x\|}), \mu(B_{x,\|X'-x\|})) \stackrel{\mathcal{L}}{=} (U, U'),$$

where U, U' are independent uniform [0, 1] random variables, we have,

$$\mu(B_{x,M}) \stackrel{\mathcal{L}}{=} \max(U,U') \stackrel{\mathcal{L}}{=} \sqrt{U} .$$

By construction, V_1 is uniform on $B_{x,||Y_2-x||}$, so that, given Y_2 ,

$$\frac{\lambda(B_{Y_2, \|Y_2 - x\|} \cup B_{Y_1, \|Y_1 - x\|})}{\lambda(B_{Y_2, \|Y_2 - x\|})} \stackrel{\mathcal{L}}{=} W ,$$

where W was defined in (1). To complete the argument, set

$$B_X = B_{X,||X-x||}, \quad B_{X'} = B_{X',||X'-x||}, \quad M = \max(||X-x||, ||X'-x||).$$

Then

$$\mu(B_X \cup B_{X'}) = \frac{\mu(B_X \cup B_{X'})}{\lambda(B_X \cup B_{X'})} \cdot \frac{\lambda(B_X \cup B_{X'})}{\lambda(B_{x,M})} \cdot \frac{\lambda(B_{x,M})}{\mu(B_{x,M})} \cdot \mu(B_{x,M})$$
$$\stackrel{\text{def}}{=} r_1 \cdot r_2 \cdot r_3 \cdot r_4.$$

Note that

$$r_1 \in [f(x)(1-\epsilon), f(x)(1+\epsilon)]$$

when $M \leq \delta$, and similarly,

$$r_3 \in \left[\frac{1}{f(x)(1+\epsilon)}, \frac{1}{f(x)(1-\epsilon)}\right]$$

when $M \leq \delta$. When (X, X') = (V, V'), we have

$$r_2 = \frac{\lambda(B_X \cup B_{X'})}{\lambda(B_{x,M})} \stackrel{\mathcal{L}}{=} W$$

with W independent of

$$r_4 = \mu(B_{x,M}) \stackrel{\mathcal{L}}{=} \sqrt{U}$$
.

Thus, since for small enough z, $\mu(B_X \cup B_{X'}) \leq z$ also $\mu(B_{x,M}) \leq z$ implies $M \leq \delta$ (as argued in the proof of (2)), for such z, we have

$$\mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, (X, X') = (V, V')\} \le \mathbb{P}\left\{\frac{1-\epsilon}{1+\epsilon}W\sqrt{U} \le z\right\} .$$

Write

$$\mathbb{P}\{\mu(B_X \cup B_{X'}) \le z\} = \mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M \le \delta\} + \mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M > \delta\}.$$

Clearly,

$$\mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M > \delta\} = 0$$

for z small enough. For such a z, we have

$$\mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M \le \delta\} = \mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M \le \delta, (X, X') \ne (V, V')\} \\
+ \mathbb{P}\{\mu(B_X \cup B_{X'}) \le z, M \le \delta, (X, X') = (V, V')\} \\
\stackrel{\text{def}}{=} q_1 + q_2.$$

We have

$$q_{1} \leq \mathbb{P}\{\mu(B_{X}) \leq z\}\mathbb{P}\{\mu(B_{X'}) \leq z\} \sup_{\rho \leq \delta} \mathbb{P}\{(X, X') \neq (V, V') \mid M \leq \rho\}$$
$$= z^{2}(1 + o(1)) \sup_{\rho \leq \delta} \mathbb{P}\{(X, X') \neq (V, V') \mid M \leq \rho\}$$
$$(\text{according to } (2))$$
$$= z^{2}(1 + o(1))\epsilon \qquad \text{(by the choice of } \delta) .$$

Also,

$$q_2 \leq \mathbb{P}\{\mu(B_X \cup B_{X'}) \leq z, M \leq \delta\}$$
$$\leq \mathbb{P}\left\{\frac{1-\epsilon}{1+\epsilon}W\sqrt{U} \leq z\right\}$$
$$\leq \left(\frac{1+\epsilon}{1-\epsilon}\right)^2 \mathbb{E}\left\{\frac{1}{W^2}\right\} z^2$$
$$= \left(\frac{1+\epsilon}{1-\epsilon}\right)^2 z^2 \alpha(d)/2 .$$

On the other hand,

$$\mathbb{P}\{\mu(B_X \cup B_{X'}) \leq z\}$$

$$\geq \mathbb{P}\{\mu(B_X \cup B_{X'}) \leq z, M \leq \delta, (X, X') = (V, V')\}$$

$$\geq \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon} \frac{\lambda(B_V \cup B_{V'})}{\lambda(B_{x,M})} \mu(B_{x,M}) \leq z, M \leq \delta, (X, X') = (V, V')\right\}$$

$$= \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon} \frac{\lambda(B_V \cup B_{V'})}{\lambda(B_{x,M})} \mu(B_{x,M}) \leq z, (X, X') = (V, V')\right\}$$

$$= A - B,$$

where

$$A = \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon}\frac{\lambda(B_V \cup B_{V'})}{\lambda(B_{x,M})}\mu(B_{x,M}) \le z\right\}$$
$$= \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon}W\sqrt{U} \le z\right\}$$
$$= \mathbb{P}\left\{U \le \left(\frac{z(1-\epsilon)}{W(1+\epsilon)}\right)^2\right\}$$
$$= \mathbb{E}\left\{\min\left\{\left(\frac{z(1-\epsilon)}{W(1+\epsilon)}\right)^2, 1\right\}\right\}$$
$$= (1-o(1))\left(\frac{1-\epsilon}{1+\epsilon}\right)^2 z^2 \frac{\alpha(d)}{2}$$

(by the dominated convergence theorem) and

$$B = \mathbb{P}\left\{\frac{1+\epsilon}{1-\epsilon}\frac{\lambda(B_V \cup B_{V'})}{\lambda(B_{x,M})}\mu(B_{x,M}) \le z, (X, X') \ne (V, V')\right\}$$
$$\le \mathbb{P}\left\{\mu(B_{x,M}) \le z, (X, X') \ne (V, V')\right\}$$
$$= \mathbb{P}\left\{\mu(B_{x,M}) \le z, M \le \delta, (X, X') \ne (V, V')\right\}$$
$$\le \mathbb{P}\left\{\mu(B_X \cup B_{X'})\frac{1-\epsilon}{1+\epsilon}\frac{1}{2^d} \le z, M \le \delta, (X, X') \ne (V, V')\right\}$$
$$= O(1)z^2\epsilon \qquad \text{(by the choice of }\delta\text{).}$$

Since ϵ was arbitrary, this concludes the proof of Part (ii) of Theorem 1.

3 Convergence in distribution

In the next result (Theorem 2) we determine the asymptotic distribution of $M_n(x)$. We do this by determining the asymptotic moments of the limiting distribution. Once again, the limit is the same for μ -almost all x.

In order to describe the asymptotic moments, for any positive integer k define the random variable

$$W_k = \frac{\lambda(B_{\overline{1},1} \cup B_{Y_1, \|Y_1\|} \cup \dots \cup B_{Y_{k-1}, \|Y_{k-1}\|})}{\lambda(B_{0,1})} ,$$

where Y_1, \ldots, Y_{k-1} are independent random variables distributed uniformly in $B_{0,1}$. Note that

$$1 = W_1 \le W_2 \le \dots \le \frac{\lambda(B_{0,2})}{\lambda(B_{0,1})} = 2^d.$$

Now we may define a non-negative random variable Z with moments

$$\mathbb{E}[Z^k] = \mathbb{E}\left[\frac{k!}{W_k^k}\right]$$

for $k \ge 1$. We may use Carleman's condition to verify that the distribution of Z is uniquely defined. Indeed, note that $\mathbb{E}[Z^k] \le k!$ and therefore

$$\sum_{k=1}^{\infty} (\mathbb{E}[Z^k])^{-1/(2k)} \ge \sum_{k=1}^{\infty} (k!)^{-1/(2k)} = \infty ,$$

and Carleman's condition is satisfied. Note that if E is an exponentially distributed random variable with mean 1, then

$$\mathbb{E}\left[E^k\right] = k! \ge \mathbb{E}[Z^k] \ge k!/2^{dk} = \mathbb{E}\left[\left(\frac{E}{2^d}\right)^k\right] .$$

We also have

$$\mathbb{E}\left[e^{sZ}\right] \le \sum_{k=0}^{\infty} s^k = \frac{1}{1-s}$$

for 0 < s < 1 and

$$\mathbb{E}\left[e^{sZ}\right] \ge \sum_{k=0}^{\infty} \left(\frac{s}{2^d}\right)^k = \frac{1}{1 - s/2^d}$$

for $0 < s < 2^d$.

The next theorem establishes the convergence announced above.

Theorem 2 Let μ have a density f. Then, for μ -almost all x, the random variable $nM_n(x)$ converges, in distribution, to Z.

Note that for the case of a Voronoi tessallation of \mathbb{R}^d defined by a Poisson point process of constant intensity, Zuyev [16] describes the distribution of the volume of the so-called "fundamental region" of the cell containing the origin, conditional on having a point at the origin, as a mixture of Gamma distributions. The fundamental region contains the Voronoi cell. Since this distribution equals the limit for the uniform density and our result is density free, the random variable Z described here is stochastically dominated by the same mixture of Gamma random variables.

In the case of d = 1 it is easily seen that Z is distributed as $(E_1 + E_2)/2$, where E_1 and E_2 are independent exponentially distributed random variables with mean 1.

Since the proof of Theorem 2 is an extension of that of part (ii) of Theorem 1, we only sketch the arguments.

Sketch of proof of Theorem 2. By the moment method, it suffices to show that for all Lebesgue points $x \in \mathbb{R}^d$ with f(x) > 0, and for all $k \ge 1$, we have

$$\mathbb{E}\left[(nM_n(x))^k\right] \to \mathbb{E}[Z^k] \; .$$

As we argued in the case k = 2 in the proof of Theorem 1,

$$\mathbb{E}\left[n^k M_n(x)^k\right] = n^k \mathbb{E}\left[(1 - Z_k(x))^{n-1}\right],$$

where

$$Z_k(x) \stackrel{\text{def}}{=} \mu(B_{X_1, \|X_1 - x\|} \cup \dots \cup B_{X_k, \|X_k - x\|})$$

is such that

$$\frac{\mathbb{P}\left\{Z_k(x) \le z\right\}}{z^k} \underset{z \to 0}{\sim} \frac{\mathbb{P}\left\{W_k U^{1/k} \le z\right\}}{z^k} ,$$

where U is uniform [0, 1], independent of W_k . Here we use the fact that

$$\max_{1 \le i \le k} \mu(B_{0, \|X_i\|}) \stackrel{\mathcal{L}}{=} \max_{1 \le i \le k} U_i \stackrel{\mathcal{L}}{=} U^{1/k}$$

with the U_i being independent and uniform on [0, 1]. The equivalence $\sim_{z\to 0}$ above can be proved by the same arguments as detailed in the proof of part (ii) of Theorem 1.

As in the proof of Theorem 1, in order to show that

$$\mathbb{E}\left[n^k M_n(x)^k\right] \to \mathbb{E}\left[\frac{k!}{W_k^k}\right] \;,$$

it suffices to show that

$$\lim_{z \downarrow 0} z^{-k} \mathbb{P}\left\{ \mu(B_{X_1, \|X_1 - x\|} \cup \dots \cup B_{X_k, \|X_k - x\|}) \le z \right\} = \mathbb{E}\left[\frac{1}{W_k^k} \right]$$

By the approximation above,

$$z^{-k} \mathbb{P}\left\{\mu(B_{X_1, \|X_1 - x\|} \cup \dots \cup B_{X_k, \|X_k - x\|}) \le z\right\} \underset{z \to 0}{\sim} z^{-k} \mathbb{P}\left\{W_k U^{1/k} \le z\right\} \underset{z \to 0}{\sim} \mathbb{E}\left[\frac{1}{W_k^k}\right] .$$

4 Some values of $\alpha(d)$

In this section we investigate the asymptotic second moment $\alpha(d)$. Since the limiting first moment equals 1, we must have that $\alpha(d) \geq 1$. On the other hand, for all d, we have $\alpha(d) \leq 2$. To see this, simply note that

$$\alpha(d) = \mathbb{E}\left[\frac{2}{W_2^2}\right] \le \mathbb{E}\left[\frac{2}{W_1^2}\right] = 2$$
.

Large values of d

Here we show that for large values of d, $\alpha(d)$ approaches 1 exponentially fast.

Theorem 3 For all $d, 1 \le \alpha(d) \le 1 + 6(3/4)^{d/2}$.

Proof. Define

$$A = B_{0,1}, \quad B = B_{\overline{1},1}, \quad \text{and} \quad C = B_{Y,||Y||},$$

where Y is uniformly distributed on A. Recall the definition of the random variable W:

$$W = \frac{\lambda(B \cup C)}{\lambda(B)}$$
 and let $U = \frac{\lambda(C)}{\lambda(B)}$

Observe that U is uniformly distributed on [0, 1]. We write

$$\begin{aligned} \alpha(d) - 1 &= \mathbb{E}\left[\frac{2}{W^2} - \frac{2}{(1+U)^2}\right] \\ &= 2\mathbb{E}\left[\frac{1}{W^2}\left(1 - \left(\frac{W}{1+U}\right)^2\right)\right] \\ &\leq 2\mathbb{E}\left[1 - \left(\frac{W}{1+U}\right)^2\right] ,\end{aligned}$$

since $W \geq 1$. We have that

$$2\left[1 - \left(\frac{W}{1+U}\right)^2\right] \le 2$$

and

$$2\left[1 - \left(\frac{W}{1+U}\right)^2\right] = 2\left[\frac{(1+U-W)(1+U+W)}{(1+U)^2}\right]$$
$$\leq 4\left[\frac{1+U-W}{1+U}\right]$$
$$= 4\left[\frac{\lambda(B\cap C)}{\lambda(B) + \lambda(C)}\right].$$

Thus

$$\alpha(d) - 1 \le 2\mathbb{E}\left[\mathbbm{1}_{Y \in B}\right] + 4\mathbb{E}\left[\frac{\lambda(B \cap C)}{\lambda(B) + \lambda(C)}\mathbbm{1}_{Y \notin B}\right] \;.$$

We finish the proof by showing that

$$\mathbb{E}\left[\mathbbm{1}_{Y\in B}\right] \le (3/4)^{d/2} \tag{4}$$

and

$$\mathbb{E}\left[\frac{\lambda(B\cap C)}{\lambda(B) + \lambda(C)}\mathbb{1}_{Y\notin B}\right] \le (3/4)^{d/2} .$$
(5)

For (4), note that

$$\mathbb{E}\left[\mathbbm{1}_{Y\in B}\right] = \frac{\lambda(A\cap B)}{\lambda(B)} \le \frac{\lambda(B_{b,\sqrt{3/4}})}{\lambda(B)} = (3/4)^{d/2} ,$$

where $b = (1/2, 0, 0, \dots 0) \in \mathbb{R}^d$, since $A \cap B \subset B_{b,\sqrt{3/4}}$ (see Figure 1.)

For (5), we bound

$$\frac{\lambda(B \cap C)}{\lambda(B) + \lambda(C)} \mathbb{1}_{Y \notin B} \leq \sup_{y \notin B} \frac{\lambda(B \cap B_{y, \|y\|})}{\lambda(B) + \lambda(B_{y, \|y\|})}$$
$$\leq \frac{\lambda(B \cap B_{a, 1})}{\lambda(B)}$$
(where $a = (1/2, \sqrt{3/4}, 0, 0, \dots, 0) \in \mathbb{R}^d$)
$$= \frac{\lambda(A \cap B)}{\lambda(B)}$$
$$\leq (3/4)^{d/2}$$



Figure 1: The sets A, B, and $B_{b,\sqrt{3/4}}$.

by arguing as above. That the supremum reached by placing Y (thus, y) at a is clear in two steps. First, the intersection $B \cap B_{y,||y||}$ can only grow by replacing y by y/||y|| since

$$B_{y,\|y\|} \subset B_{y/\|y\|,1}.$$

Next, of all the points on the surface of $B_{\overline{1},1}$, but outside $B_{\overline{1},1}$, the intersection $\lambda(B \cap B_{y/||y||,1})$ is maximized by placing y at a.

Small values of d

It is not difficult to see that $\alpha(1) = 3/2$. Indeed, according to (1),

$$\overline{B} = B_{\overline{1},1} \cup B_{Y,||Y||} = B_{1,1} \cup B_{Y,|Y|} = \begin{cases} B_{1,1} & \text{if } Y \ge 0\\ B_{1,1} \cup B_{Y,|Y|} & \text{if } Y < 0 \end{cases},$$

we have

$$W = \frac{\lambda(\overline{B})}{\lambda(B_{0,1})} \stackrel{\mathcal{L}}{=} \left\{ \begin{array}{ll} 1 & \text{ with probability } 1/2, \\ 1+U & \text{ with probability } 1/2, \end{array} \right.$$

where U is uniform [0, 1]. Hence,

$$\alpha(1) = \mathbb{E}\left[\frac{2}{W^2}\right] = \frac{1}{2}\left(\frac{2}{1} + \mathbb{E}\left[\frac{2}{(1+U)^2}\right]\right) = 1 + \mathbb{E}\left[\frac{1}{(1+U)^2}\right] = 3/2 .$$

Previous work has considered the variance of the Lebesgue measure of the Voronoi cell containing the origin defined by a homogeneous Poisson process, conditioned on the fact that a point falls in the origin. From these results, we deduce values of $\alpha(d)$ for d = 2, 3. Indeed Gilbert [7], Brakke [3], and Hayen and Quine [8] showed that $\alpha(2) \approx 1.2801760409267$ while Gilbert [7] and Brakke [4] showed $\alpha(3) \approx 1.179032437845$.

5 On the diameter of a typical cell

Here we prove that, independently of the density, for μ -almost all $x \in \mathbb{R}^d$, conditional on $X_1 = x$, the diameter of the Voronoi cell centered at X_1 converges to zero, in probability, at a rate of $n^{-1/d}$ in the sense specified in Theorem 4. More precisely, for any $x \in \mathbb{R}^d$, let $D_n(x)$ be the diameter of the Voronoi cell S_1 with nucleus x associated with the point process $\{x, X_2, \ldots, X_n\}$. Then we have the following:

Theorem 4 Let μ have a density f. Then for μ -almost all $x \in \mathbb{R}^d$,

$$\lim_{t \to \infty} \limsup_{n \to \infty} \mathbb{P}\left\{ n^{1/d} D_n(x) \ge t \right\} = 0 \; .$$

Proof. A set $C \subset \mathbb{R}^d$ is a cone of angle $\theta \in (0, \pi/2)$ centered at 0 if there exists an $x \in \mathbb{R}^d$ with ||x|| = 1 such that

$$C = \left\{ y \in \mathbb{R}^d : \frac{(x, y)}{\|y\|} \ge \cos(\theta/2) \right\}$$

Let γ_d be the minimal number of cones $C_1, \ldots, C_{\gamma_d}$ of angle $\pi/4$ centered at 0 such that their union covers \mathbb{R}^d . (For example, $\gamma_2 = 8$.) Let $R_{n,j}$, $j = 1, \ldots, \gamma_d$, be the distance between x and the nearest neighbor among X_2, \ldots, X_n belonging to $x + C_j$ (i.e., the cone C_j translated by x). Define $R_{n,j} = \infty$ if no such point exists.

We bound the diameter of the Voronoi cell S_1 by observing that

$$D_n(x) \leq \sqrt{2} \max_{j=1,\dots,\gamma_d} R_{n,j}$$
.

To see this, consider an arbitrary point $y \in S_1$ and let $x + C_j$ be a cone (among the covering cones) containing y. If ξ_j is the nearest neighbor of x among X_2, \ldots, X_n belonging to $x + C_j$, then $||y - x|| \le ||y - \xi_j||$ and therefore $||y - x|| \le R_{n,j}/(2\cos(\pi/4))$. Thus, the diameter of S_1 is at most $\max_{j=1,\ldots,\gamma_d} R_{n,j}/\cos(\pi/4)$.

A simple extension of the Lebesgue density theorem implies that if $A = B_{0,1}$ is the unit ball centered at the origin, then for μ -almost all $x \in \mathbb{R}^d$,

$$\min_{j=1,\dots,\gamma_d} \frac{\int_{x+r[C_j\cap A]} f}{\lambda(r[C_j\cap A])} \to f(x) \quad \text{as} \quad r \downarrow 0 ,$$
(6)

where $r[C_j \cap A] = \{rx : x \in C_j \cap A\}$. Thus, for μ -almost all x, there exists R(x) > 0 such that for all $0 < r \le R(x)$,

$$\min_{j=1,\dots,\gamma_d} \int_{x+r[C_j \cap A]} f \ge r^d \frac{f(x)}{2} \lambda(C_1 \cap A) .$$

If f(x) = 0 or x does not satisfy (6), set R(x) = 0. For any t > 0, we have

$$\left\{ D_n(x) > tn^{-1/d} \right\} \subset \left\{ \max_{j=1,\dots,\gamma_d} R_{n,j} > \frac{tn^{-1/d}}{\sqrt{2}} \right\}$$
$$\subset \bigcup_{j=1}^{\gamma_d} \left\{ x + \frac{tn^{-1/d}}{\sqrt{2}} [C_j \cap A] \text{ has no point among } X_2, \dots, X_n \right\} .$$

Thus, we have

$$\mathbb{P}\left\{n^{1/d}D_n(x) \ge t\right\}$$

$$\leq \sum_{j=1}^{\gamma_d} \mathbb{P}\left\{x + \frac{tn^{-1/d}}{\sqrt{2}}[C_j \cap A] \text{ has no point among } X_2, \dots, X_n\right\}.$$

We bound the probability of each event in the union as follows.

$$\mathbb{P}\left\{x + \frac{tn^{-1/d}}{\sqrt{2}}[C_j \cap A] \text{ has no point among } X_2, \dots, X_n\right\} \\
\leq \mathbb{P}\left\{x + \min\left(R(x), \frac{tn^{-1/d}}{\sqrt{2}}\right)[C_j \cap A] \text{ has no point among } X_2, \dots, X_n\right\} \\
= \left(1 - \mu\left(x + \min\left(R(x), \frac{tn^{-1/d}}{\sqrt{2}}\right)[C_j \cap A]\right)\right)^{n-1} \\
\leq \left(1 - \left(\min\left(R(x), \frac{tn^{-1/d}}{\sqrt{2}}\right)\right)^d \lambda(C_1 \cap A) \frac{f(x)}{2}\right)^{n-1} \\
\leq \exp\left(-(n-1)\min\left(R(x)^d, \frac{t^d n^{-1}}{\sqrt{2^d}}\right)\left(\lambda(C_1 \cap A) \frac{f(x)}{2}\right)\right)$$

and the theorem follows since $R(x)^d f(x) > 0$ for μ -almost all x.

Note that it follows from the proof that, for μ -almost every x, there exists a positive constant C(x) such that

$$\mathbb{P}\left\{n^{1/d}D_n(x) \ge t\right\} \le \gamma_d \exp(-(n-1)C(x))$$

and therefore the diameter $D_n(x)$ has an exponentially bounded tail. This phenomenon has been observed before for the Poisson-Voronoi tessellation, see, e.g., Maier, Mayer, and Schmidt [11].

Acknowledgements. We thank the associate editor and a referee for extremely useful feedback that helped simplify some proofs and significantly improve presentation.

References

- F. Baccelli, M. Klein, M. Lebourges and S. A. Zuyev, Stochastic geometry and architecture of communication networks. *Telecommunications Systems*, 7:209–227, 1997.
- [2] G. Biau, L. Devroye, Lectures on the Nearest Neighbor Method. Springer-Verlag, New York, 2015.
- [3] K. A. Brakke, Statistics of random plane Voronoi tessellations, Dept. of Math. Sciences, Susquehanna University, Selinsgrove (Pennsylvania), 1–30, 1985. http://www.susqu.edu/brakke/aux/downloads/papers/vorplane.pdf.
- [4] K. A. Brakke, Statistics of three dimensional random Voronoi tessellations, Dept. of Math. Sciences, Susquehanna University, Selinsgrove (Pennsylvania), 1–17, 1985. http://www.susqu.edu/brakke/aux/downloads/papers/3d.pdf.
- [5] S.N. Chiu, D. Stoyan, W.S. Kendall, and J. Mecke, *Stochastic Geometry and its Applications*. Third edition. Akademie-Verlag, Berlin, 2013.
- [6] L. Devroye and L. Györfi, Nonparametric Density Estimation: The L_1 View. John Wiley, New York, 1985.
- [7] E. N. Gilbert, Random subdivisions of space into crystals. Ann. Math. Statist., 33:958-972, 1962.
- [8] A. Hayen and M. P. Quine, Areas of components of a Voronoi polygon in a homogeneous Poisson process in the plane, Adv. Appl. Prob., 34:281-291, 2002.

- [9] L. Heinrich, R. Körner, N. Mehlhorn and L. Muche, Numerical and analytical computation of some second-order characteristics of spatial PoissonVoronoi tessellations, *Statistics*, 31:235-259, 1998.
- [10] L. Heinrich and L. Muche, Second-order properties of the point process of nodes in a stationary Voronoi tessellation, *Math. Nachr.*, 281:350-375, 2008.
- [11] R. Maier, J. Mayer, and V. Schmidt, Distributional properties of the typical cell of stationary iterated tessellations, *Mathematical Methods of Operations Research*, 59:287– 302, 2004.
- [12] J. Møller, Lectures on Random Voronoi Tessellations, Lecture Notes in Statistics Vol. 87, Springer, New York, 1994.
- [13] J. Møller and D. Stoyan, Stochastic geometry and random tessellations. In *Tessellations in the Sciences: Virtues, Techniques and Applications of Geometric Tilings*, eds. R. van de Weijgaert, G. Vegter, V. Icke, and J. Ritzerveld. Springer Verlag, 2007.
- [14] A. Okabe, B. Boots, K. Sugihara, and S.N. Chiu, Spatial Tessellations: Concepts and Applications of Voronoi Diagrams. Second edition. John Wiley, Chichester, England, 2000.
- [15] R. Wheeden and A. Zygmund, *Measure and Integral*. Marcel Dekker, New York, 1977.
- [16] S. A. Zuyev, Estimates for distributions of the Voronoi polygon's geometric characteristics, Random Structures and Algorithms, 3:149-162, 1992.

Luc Devroye McGill University **E-Mail:** lucdevroye@gmail.com

László Györfi Budapest University of Technology and Economics **E-Mail:** gyorfi@cs.bme.hu

Gábor Lugosi ICREA and Pompeu Fabra University **E-Mail:** gabor.lugosi@upf.edu

Harro Walk Universität Stuttgart **E-Mail:** harro.walk@t-online.de

Erschienene Preprints ab Nummer 2012-001

Komplette Liste: http://www.mathematik.uni-stuttgart.de/preprints

- 2016-006 Devroye, L.; Györfi, L.; Lugosi, G.; Walk, H.: On the measure of Voronoi cells
- 2016-005 Kohls, C.; Kreuzer, C.; Rösch, A.; Siebert, K.G.: Convergence of Adaptive Finite Elements for Optimal Control Problems with Control Constraints
- 2016-004 *Blaschzyk, I.; Steinwart, I.:* Improved Classification Rates under Refined Margin Conditions
- 2016-003 *Feistauer, M.; Roskovec, F.; Sändig, AM.:* Discontinuous Galerkin Method for an Elliptic Problem with Nonlinear Newton Boundary Conditions in a Polygon
- 2016-002 *Steinwart, I.:* A Short Note on the Comparison of Interpolation Widths, Entropy Numbers, and Kolmogorov Widths
- 2016-001 Köster, I.: Sylow Numbers in Spectral Tables
- 2015-016 Hang, H.; Steinwart, I.: A Bernstein-type Inequality for Some Mixing Processes and Dynamical Systems with an Application to Learning
- 2015-015 *Steinwart, I.:* Representation of Quasi-Monotone Functionals by Families of Separating Hyperplanes
- 2015-014 Muhammad, F.; Steinwart, I.: An SVM-like Approach for Expectile Regression
- 2015-013 Nava-Yazdani, E.: Splines and geometric mean for data in geodesic spaces
- 2015-012 Kimmerle, W.; Köster, I.: Sylow Numbers from Character Tables and Group Rings
- 2015-011 *Györfi, L.; Walk, H.:* On the asymptotic normality of an estimate of a regression functional
- 2015-010 Gorodski, C, Kollross, A.: Some remarks on polar actions
- 2015-009 Apprich, C.; Höllig, K.; Hörner, J.; Reif, U.: Collocation with WEB-Splines
- 2015-008 *Kabil, B.; Rodrigues, M.:* Spectral Validation of the Whitham Equations for Periodic Waves of Lattice Dynamical Systems
- 2015-007 Kollross, A.: Hyperpolar actions on reducible symmetric spaces
- 2015-006 *Schmid, J.; Griesemer, M.:* Well-posedness of Non-autonomous Linear Evolution Equations in Uniformly Convex Spaces
- 2015-005 *Hinrichs, A.; Markhasin, L.; Oettershagen, J.; Ullrich, T.:* Optimal quasi-Monte Carlo rules on higher order digital nets for the numerical integration of multivariate periodic functions
- 2015-004 *Kutter, M.; Rohde, C.; Sändig, A.-M.:* Well-Posedness of a Two Scale Model for Liquid Phase Epitaxy with Elasticity
- 2015-003 *Rossi, E.; Schleper, V.:* Convergence of a numerical scheme for a mixed hyperbolic-parabolic system in two space dimensions
- 2015-002 *Döring, M.; Györfi, L.; Walk, H.:* Exact rate of convergence of kernel-based classification rule
- 2015-001 *Kohler, M.; Müller, F.; Walk, H.:* Estimation of a regression function corresponding to latent variables
- 2014-021 *Neusser, J.; Rohde, C.; Schleper, V.:* Relaxed Navier-Stokes-Korteweg Equations for Compressible Two-Phase Flow with Phase Transition
- 2014-020 *Kabil, B.; Rohde, C.:* Persistence of undercompressive phase boundaries for isothermal Euler equations including configurational forces and surface tension
- 2014-019 *Bilyk, D.; Markhasin, L.:* BMO and exponential Orlicz space estimates of the discrepancy function in arbitrary dimension

- 2014-018 *Schmid, J.:* Well-posedness of non-autonomous linear evolution equations for generators whose commutators are scalar
- 2014-017 *Margolis, L.:* A Sylow theorem for the integral group ring of PSL(2,q)
- 2014-016 *Rybak, I.; Magiera, J.; Helmig, R.; Rohde, C.:* Multirate time integration for coupled saturated/unsaturated porous medium and free flow systems
- 2014-015 *Gaspoz, F.D.; Heine, C.-J.; Siebert, K.G.:* Optimal Grading of the Newest Vertex Bisection and *H*¹-Stability of the *L*₂-Projection
- 2014-014 Kohler, M.; Krzyżak, A.; Walk, H.: Nonparametric recursive quantile estimation
- 2014-013 Kohler, M.; Krzyżak, A.; Tent, R.; Walk, H.: Nonparametric quantile estimation using importance sampling
- 2014-012 *Györfi, L.; Ottucsák, G.; Walk, H.:* The growth optimal investment strategy is secure, too.
- 2014-011 Györfi, L.; Walk, H.: Strongly consistent detection for nonparametric hypotheses
- 2014-010 *Köster, I.:* Finite Groups with Sylow numbers $\{q^x, a, b\}$
- 2014-009 Kahnert, D.: Hausdorff Dimension of Rings
- 2014-008 Steinwart, I.: Measuring the Capacity of Sets of Functions in the Analysis of ERM
- 2014-007 *Steinwart, I.:* Convergence Types and Rates in Generic Karhunen-Loève Expansions with Applications to Sample Path Properties
- 2014-006 Steinwart, I.; Pasin, C.; Williamson, R.; Zhang, S.: Elicitation and Identification of Properties
- 2014-005 *Schmid, J.; Griesemer, M.:* Integration of Non-Autonomous Linear Evolution Equations
- 2014-004 *Markhasin, L.:* L_2 and $S_{p,q}^r B$ -discrepancy of (order 2) digital nets
- 2014-003 *Markhasin, L.:* Discrepancy and integration in function spaces with dominating mixed smoothness
- 2014-002 Eberts, M.; Steinwart, I.: Optimal Learning Rates for Localized SVMs
- 2014-001 *Giesselmann, J.:* A relative entropy approach to convergence of a low order approximation to a nonlinear elasticity model with viscosity and capillarity
- 2013-016 Steinwart, I.: Fully Adaptive Density-Based Clustering
- 2013-015 *Steinwart, I.:* Some Remarks on the Statistical Analysis of SVMs and Related Methods
- 2013-014 *Rohde, C.; Zeiler, C.:* A Relaxation Riemann Solver for Compressible Two-Phase Flow with Phase Transition and Surface Tension
- 2013-013 Moroianu, A.; Semmelmann, U.: Generalized Killling spinors on Einstein manifolds
- 2013-012 Moroianu, A.; Semmelmann, U.: Generalized Killing Spinors on Spheres
- 2013-011 Kohls, K; Rösch, A.; Siebert, K.G.: Convergence of Adaptive Finite Elements for Control Constrained Optimal Control Problems
- 2013-010 *Corli, A.; Rohde, C.; Schleper, V.:* Parabolic Approximations of Diffusive-Dispersive Equations
- 2013-009 Nava-Yazdani, E.; Polthier, K.: De Casteljau's Algorithm on Manifolds
- 2013-008 *Bächle, A.; Margolis, L.:* Rational conjugacy of torsion units in integral group rings of non-solvable groups
- 2013-007 Knarr, N.; Stroppel, M.J.: Heisenberg groups over composition algebras
- 2013-006 Knarr, N.; Stroppel, M.J.: Heisenberg groups, semifields, and translation planes

- 2013-005 *Eck, C.; Kutter, M.; Sändig, A.-M.; Rohde, C.:* A Two Scale Model for Liquid Phase Epitaxy with Elasticity: An Iterative Procedure
- 2013-004 Griesemer, M.; Wellig, D.: The Strong-Coupling Polaron in Electromagnetic Fields
- 2013-003 *Kabil, B.; Rohde, C.:* The Influence of Surface Tension and Configurational Forces on the Stability of Liquid-Vapor Interfaces
- 2013-002 Devroye, L.; Ferrario, P.G.; Györfi, L.; Walk, H.: Strong universal consistent estimate of the minimum mean squared error
- 2013-001 *Kohls, K.; Rösch, A.; Siebert, K.G.:* A Posteriori Error Analysis of Optimal Control Problems with Control Constraints
- 2012-013 *Diaz Ramos, J.C.; Dominguez Vázquez, M.; Kollross, A.:* Polar actions on complex hyperbolic spaces
- 2012-012 Moroianu; A.; Semmelmann, U.: Weakly complex homogeneous spaces
- 2012-011 Moroianu; A.; Semmelmann, U.: Invariant four-forms and symmetric pairs
- 2012-010 Hamilton, M.J.D.: The closure of the symplectic cone of elliptic surfaces
- 2012-009 Hamilton, M.J.D.: Iterated fibre sums of algebraic Lefschetz fibrations
- 2012-008 Hamilton, M.J.D.: The minimal genus problem for elliptic surfaces
- 2012-007 *Ferrario, P.:* Partitioning estimation of local variance based on nearest neighbors under censoring
- 2012-006 Stroppel, M.: Buttons, Holes and Loops of String: Lacing the Doily
- 2012-005 Hantsch, F.: Existence of Minimizers in Restricted Hartree-Fock Theory
- 2012-004 Grundhöfer, T.; Stroppel, M.; Van Maldeghem, H.: Unitals admitting all translations
- 2012-003 Hamilton, M.J.D.: Representing homology classes by symplectic surfaces
- 2012-002 Hamilton, M.J.D.: On certain exotic 4-manifolds of Akhmedov and Park
- 2012-001 Jentsch, T.: Parallel submanifolds of the real 2-Grassmannian